



Health risk assessment related to pinnatoxins in French shellfish

Nathalie Arnich, Eric Abadie, Nicolas Delcourt, Valérie Fessard, Jean-Marc Fremy, Vincent Hort, Emmeline Lagrange, Thomas Maignien, Jordi Molgó, Marie-Bénédicte Peyrat, et al.

► To cite this version:

Nathalie Arnich, Eric Abadie, Nicolas Delcourt, Valérie Fessard, Jean-Marc Fremy, et al.. Health risk assessment related to pinnatoxins in French shellfish. *Toxicon*, 2020, 180, pp.1-10. 10.1016/j.toxicon.2020.03.007 . anses-02861821

HAL Id: anses-02861821

<https://anses.hal.science/anses-02861821>

Submitted on 20 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Type of paper: Full-Length Research Papers

Health risk assessment related to pinnatoxins in French shellfish

Nathalie ARNICH^{1a}, Eric ABADIE², Nicolas DELCOURT³, Valérie FESSARD^{1b}, Jean-Marc FREMY¹,
Vincent HORT^{1c}, Emmeline LAGRANGE⁴, Thomas MAIGNIEN^{1a}, Jordi MOLGÓ⁵, Marie-Bénédicte
PEYRAT^{1a}, Jean-Paul VERNOUX⁶, César MATTEI⁷

¹ANSES (French Agency for Food, Environmental and Occupational Health and Safety), ^aRisk
Assessment Directorate, Maisons-Alfort, France, corresponding author: nathalie.arnich@anses.fr;

^bToxicology of Contaminants Unit, Fougères, France ; ^cLaboratory for Food Safety, Maisons-Alfort,
France

²IFREMER (French Research Institute for Exploitation of the Sea), Centre for Marine Biodiversity,
Exploitation and Conservation (MARBEC), Sète, France

³Poison Control Centre, Toulouse-Purpan University Hospital and Toulouse NeuroImaging Centre
(ToNIC), Toulouse, France

⁴Department of Neurology, Reference Center of Neuromuscular Disease, Grenoble University
Hospital, France

⁵CEA (French Alternative Energies and Atomic Energy Commission), CNRS, Gif-sur-Yvette, France

⁶Normandie University, Caen, France

⁷Mitochondrial and Cardiovascular Pathophysiology, UMR CNRS 6015, INSERM U1083, Angers
University, Angers, France

ABSTRACT

Pinnatoxins (PnTXs) are a group of emerging marine biotoxins produced by the benthic dinoflagellate *Vulcanodinium rugosum*, currently not regulated in Europe or in any other country in the world. In France, PnTXs were detected for the first time in 2011, in mussels from the Ingril lagoon (South of France, Mediterranean coast). Since then, analyses carried out in mussels from this lagoon have shown high concentrations of PnTXs for several months each year. PnTXs have also been detected, to a lesser extent, in mussels from other Mediterranean lagoons and on the Atlantic and Corsican coasts. In the French data, the main analog is PnTX G (low levels of PnTX A are also present in some samples). No cases of PnTXs poisoning in humans have been reported so far in France or anywhere else in the world. In mice, PnTXs induce acute neurotoxic effects, within a few minutes after oral administration. Clinical signs of toxicity include decreased mobility, paralysis of the hind legs, tremors, jumps and breathing difficulties leading to death by respiratory arrest at high doses. The French agency for food safety (ANSES) recently conducted a review of the state of knowledge related to PnTXs and *V. rugosum*. Based on (i) the clinical signs of toxicity in mice, (ii) the mode of action of PnTXs as nicotinic acetylcholine receptor competitive antagonists and (iii) knowledge on drugs and natural toxins with PnTX-related pharmacology, potential human symptoms have been extrapolated and proposed. In this work, a provisional acute benchmark value for PnTX G of 0.13 µg/kg bw per day has been derived from an oral acute toxicity study in mice. Based on this value and a large shellfish meat portion size of 400g, a concentration lower than 23 µg PnTX G/kg shellfish meat is not expected to result in adverse effects in humans. ANSES recommends taking into account PnTXs in the French official monitoring program for shellfish production and identified data gaps to refine health risk assessment.

KEYWORDS: Pinnatoxins, shellfish, emerging marine biotoxins, risk assessment

1. Introduction

Pinnatoxins (PnTXs) belong to the group of cyclic imines considered as emerging marine biotoxins (Efsa, 2010a), which, to date, includes 40 compounds, without considering acyl esters produced by shellfish metabolism. This group includes different families determined by their structural characteristics: prorocentrolides, spiroporocentrimine, gymnodimines (GYMs), spirolides (SPXs), pinnatoxins (PnTXs), pteriatoxins (PtTXs) and portimine. To date, 8 PnTXs (named A to H) have been identified (reviewed by Molgó et al., 2017). PnTXs are soluble in solvents such as acetone, isopropanol and methanol (Zendong et al. 2014). They are amphoteric compounds, i.e. with both acidic and basic properties, which explains their relative water solubility. Due to their lipophilic properties, they can be detected during the mouse bioassay used to screen for lipophilic marine biotoxins.

Vulcanodinium rugosum is the producer of PnTXs identified by Nézan and Chomerat (2011), based on water samples from a Mediterranean lagoon (Ingril, France). *V. rugosum* also produce portimine (Selwood et al., 2013; Abadie et al., 2016). This benthic dinoflagellate was a new species belonging to a new genus. No other producer is reported in the literature. The identification of this dinoflagellate in France originates from an atypical situation that occurred in 2006, as part of official monitoring of shellfish production areas. The mouse bioassay used to screen for lipophilic marine biotoxins had revealed unusual neurotoxic effects after the injection of extracts from mussels of the Ingril lagoon. In fact, the symptoms observed for lipophilic toxins are gastrointestinal (diarrhea), and not neurotoxic. Such neurotoxicity could not be related to the presence of regulated marine biotoxins (Amnesic, Paralytic, Diarrhetic toxins) screened for by chemical analysis. In addition, observations of water samples did not conclude to the presence of any microalgae species known to produce neurotoxins. *Vulcanodinium rugosum* has also been reported in New Zealand (Rhodes et al., 2010), Australia (Rhodes et al., 2011; Munday et al., 2012), Japan (Smith et al., 2011), China Sea (Zeng et al., 2012), Mexico (Hernandez-Becerril et al., 2013), Arabian Gulf (Al Muftah et al., 2016) and Cuba (Moreira-Gonzalez et al., 2018).

PnTXs in shellfish are mainly analysed by physico-chemical methods using liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS). Biological (mouse bioassay) or biochemical (functional tests) methods can also be used. None of the methods has yet undergone inter-laboratory

validation or standardisation. In addition, there are very few standards (reference substances) available for PnTXs, which limits and complicates their detection and quantification. Indeed, to date, only PnTX A and G are marketed as calibration solutions and only PnTX G (Figure 1) has been certified.

Since the first identification of *V. rugosum* in France, high concentrations of PnTXs have been measured in mussels from the Ingril lagoon for several months each year. In 2012, the French Research Institute for Exploitation of the Sea (Ifremer) reported that PnTXs concentrations varied greatly depending on the years (2010, 2011 and 2012), with a maximum of 1244 µg of PnTX G per kg of shellfish (wet weight) in 2010 (Ifremer, 2012). However, *V. rugosum* is rarely detected in water samples collected for the phytoplankton surveillance (in the water column), which can be explained by the benthic nature of this dinoflagellate.

The aims of this work were 1) to review the toxicity data of PnTXs to derive an acute oral health-based guidance value, 2) estimate exposure of shellfish consumers to PnTXs, 3) identify if there might be a public health concern regarding the levels of contamination reported in certain French shellfish production areas, and 4) provide recommendations for the monitoring PnTXs in the marine environment.




Figure 1: Structure of PnTX G (modified from Araoz et al., 2011).

2. *Vulcanodinium rugosum*

As almost all dinoflagellates, *V. rugosum* (Figure 2) is assumed to have two distinct life phases: a vegetative propagation phase (asexual reproduction) and a sexual phase (Zeng et al., 2012). The asexual phase probably corresponds to the pelagic phase, when the cells are present in the water

column. The sexual reproduction corresponds to a benthic phase, which could result in the formation of a planozygote that produces a resistance cyst. The pelagic and benthic phases are closely related, but the environmental and/or physiological factors controlling the transition from one phase to the other are not yet known.

The formation of resistance cysts, which can occur at the end of sexual reproduction, is a crucial step for survival in adverse environmental conditions due to chromosomal mixing, which increases intra-species genetic diversity. Cysts are also one of the ways in which the species spreads, through the transfer of sediments or shellfish from one area to another (cysts are present in the gastrointestinal tract and in intervalvular liquid), as well as via ship ballast water (Garrett et al., 2014). Therefore, identification of the resistance forms and knowledge of their distribution area are important for preventing expansion and contamination. Similarly, determination of the key factors for growth and toxin production by *V. rugosum* is essential for better understanding the risk associated with this new species.

Temperature is the most important factor for the growth of *V. rugosum*. The data suggest that *V. rugosum* is a thermophilic species, consistent with its development in the Ingril lagoon from June to September and the highest concentrations of PnTX G found in mussels during this period (Abadie et al., 2016).




Figure 2: Micrograph of a cell of *Vulcanodinium rugosum*

3. Hazard characterisation

3.1 Acute *in vivo* toxicity with purified PnTXs

In vivo acute toxicity studies with purified toxins were reviewed using an analysis grid and assessed for quality using ToxRTool (Toxicological data Reliability assessment Tool, <https://eurl-ecvam.jrc.ec.europa.eu/about-ecvam/archivepublications/toxrtool>), a tool that ranks the studies according to the Klimisch rating (Klimish et al., 1997).

Data on the acute toxicity of purified PnTXs are very limited. Indeed, the available studies were carried out in only one species (mice), one sex (females, which are considered to be more sensitive than males by OECD guidelines) and with very few animals tested per dose (1 to 8 mice, when the number is specified). The results are presented in table 1.

Orally, only three studies are available and focused on PnTX E, F, G and H (Munday *et al.*, 2012; Selwood *et al.*, 2014; ANSES-University of Trieste-CNRS report, 2014 also published in Sosa *et al.*, 2020). We did not find any information on the toxicity by the oral route for PtTXs (metabolites of certain PnTXs) and portimine (a toxin also produced by *V. rugosum*). Regarding the intraperitoneal (IP) route, four studies have been published on PnTX E, F, G and H and portimine (Munday *et al.*, 2012; Selwood *et al.*, 2010, 2013, 2014). We identified additional unpublished information regarding PnTX A (personal communication from J. Molgó), but did not find any robust information regarding PtTXs.

Despite the limited number of studies for each PnTX and the low number of mice tested per dose, a consistent set of information enables outlining the main characteristics of the acute toxicity of this toxin group. Firstly, the toxicity of the PnTXs is rapid, with symptoms appearing within minutes of administration (whether oral or IP). This fact was already known, since PnTXs (like other cyclic imines) belong to the fast-acting toxins group (Efsa, 2010a). The second point relies on neurotoxic symptoms, quickly leading to the mouse's death by respiratory arrest. Clinical signs of toxicity, regardless of the route of administration and the PnTX analogue, include decreased mobility (sometimes preceded by an initial phase of hyperactivity immediately following administration), paralysis of the hind legs and breathing difficulties (Munday *et al.*, 2012), with tremors and jumps also being reported (ANSES-University of Trieste-CNRS Report, 2014; Sosa *et al.*, 2020). In the study by Munday *et al.* (2012), the authors reported that at sublethal doses (without specifying which ones), some mice recovered completely after exhibiting symptoms.

LD50 values vary according to the PnTX analogue and route of administration (Table 1). Orally, LD50 values range from 25 to 2,800 µg/kg bw (for PnTX F and PnTX E respectively). The analogues tested can be ranked as follows in decreasing order of toxicity: PnTX F > PnTX G ~ PnTX H >> PnTX E. By

IP route, LD50 range from 13 to 115 µg/kg bw (for PnTX F and PnTX A respectively) and the analogues can be ranked as follows: PnTX F > PnTX G > PnTX E > PnTX H > PnTX A.

LD99 values for some PnTXs and PtTXs have been reported in the literature, but these data were considered not sufficiently reliable due to a lack of information on the protocol used (Uemura *et al.*, 1995; Chou *et al.*, 1996; McCauley *et al.*, 1998; Takada *et al.*, 2001).

Portimine has lower acute IP toxicity than that of PnTXs, with an estimated LD50 of 1,570 µg/kg bw. No effect was observed at 500 and 700 µg/kg bw (Selwood *et al.*, 2013). The authors indicated that the signs of toxicity prior to death in mice appeared less rapidly after IP administration compared to PnTXs (without however mentioning which signs of toxicity were observed). Toxicity of portimine by oral administration is unknown.

Table 1: Acute *in vivo* toxicity in mice of PnTXs and portimine

1a. Oral administration

Toxin (purity)	Route of administration and number of mice	LD50 (µg/kg bw)	MTD (µg/kg bw)	References	Study quality
PnTX E*	Gavage Number of mice not specified (OECD GL 425) Fed (non-fasted) mice	2800 CI95: 2380-3000	600	Munday <i>et al.</i> , 2012	ToxRTool: 13 Klimisch: 2
PnTX F*	Gavage Number of mice not specified (OECD GL 425) Fed (non-fasted) mice	25 CI95: 19.1-35.1	9.9	Munday <i>et al.</i> , 2012	ToxRTool: 13 Klimisch: 2
PnTX F*	16h fasted mice	29.9 CI95: 25-32	Not determined		
	In cream cheese Number of animals not specified (OECD GL 425) Fed (non-fasted) mice	50 CI95: 39.4-62.8	16.0	Munday <i>et al.</i> , 2012	ToxRTool: 13 Klimisch: 2
	In peanut butter Number of mice not specified (OECD GL 425) Fed (non-fasted) mice	50 CI95: 37.9-71.5	Not determined		
	In dry mouse food Number of mice not	50 CI95: 37.9-71.5	Not determined		

Toxin (purity)	Route of administration and number of mice	LD50 (µg/kg bw)	MTD (µg/kg bw)	References	Study quality
	specified (OECD GL 425) 16h fasted mice In cream cheese, 16h fasted mice In peanut butter, 16h fasted mice	77 CI95: not calculated 50 CI95: 39.4-62.8	Not determined Not determined		
PnTX G*	Gavage Number of mice not specified (OECD GL 425) Fed (non-fasted) mice	150 CI95: 105-100	75	Munday <i>et al.</i> , 2012	ToxRTool: 13 Klimisch: 2
PnTX G (declared purity 100%)	Gavage Groups of 3 to 8 mice (8 for the control group; 3 for the doses 8, 20, 50, 120 µg/kg bw; 5 for the doses 220, 300, 370, 400 µg/kg bw) Mice fasted for 3 hours before administration	208 CI95: 155-281	120	ANSES-University of Trieste-CNRS report, 2014; Sosa <i>et al.</i> , 2020	ToxRTool: 17 Klimisch: 2
PnTX G*	In cream cheese Number of mice not specified (OECD GL 425) Fed (non-fasted) mice	400 CI95: 380-470	153	Munday <i>et al.</i> , 2012	ToxRTool: 13 Klimisch: 2
PnTX H*	Gavage Number of animals not specified Fasting not specified	163 CI95: 139-175		Selwood <i>et al.</i> , 2014	ToxRTool: 4 This low score is due to the fact that a lot of details are missing in the article, but this team has already described the protocol for other PnTXs

* Purity verified by NMR according to the authors but percentage not mentioned in the publication

CI95: 95% confidence interval

MTD (Maximum Tolerated Dose): dose at which no mortality or clinical signs are observed

1b. Intraperitoneal (IP) administration

Toxin (purity)	Route of administration and number of mice	LD50 (µg/kg bw)	MTD (µg/kg bw)	References	Study quality
Synthetic PnTX A (purity > 97%)	IP n=18 mice, 9 doses tested (1 to 3 mice/dose)	114.8		J. Molgó (personal communication)	ToxRTool: 16

Toxin (purity)	Route of administration and number of mice	LD50 (µg/kg bw)	MTD (µg/kg bw)	References	Study quality
	Fed (non-fasted) mice				
PnTX E*	IP Number of animals not specified (OECD GL 425)	57 (fed) CI95: 39.7-75.3 48 (16h fasted) CI95: 33.5-63.5	22 Not determined	Munday <i>et al.</i> , 2012	ToxRTool: 13
	IP (OECD GL 425) n=2 at 36 µg/kg, n=3 at 45 µg/kg, n=1 at 54 µg/kg, n=1 at 60 µg/kg.	45 (fed) CI95: 32-58		Selwood <i>et al.</i> , 2010	ToxRTool: 15
PnTX F*	IP Number of animals not specified (OECD GL 425)	12.7 (fed) CI95: 9.5-14.6 14.9 (16h fasted) CI95: 12.6-15.8	3.2 Not determined	Munday <i>et al.</i> , 2012	ToxRTool: 13
	IP (OECD GL 425) n=1 at 10.1 µg/kg, n=1 at 12.7 µg/kg, n=3 at 16.0 µg/kg, n=2 at 20.1 µg/kg.	16 (fed) CI95: 12-23		Selwood <i>et al.</i> , 2010	ToxRTool: 15
PnTX G*	IP Number of animals not specified (OECD GL 425)	48 (fed) CI95: 36.3-68.1 42.7 (16h fasted) CI95: 40-50	18.8 Not determined	Munday <i>et al.</i> , 2012	ToxRTool: 13
PnTX G*	IP (OECD GL 425) n=2 at 40 µg/kg, n=3 at 50 µg/kg, n=1 at 60 µg/kg.	50 (fed) CI95: 35-66		Selwood <i>et al.</i> , 2010	ToxRTool: 15
Synthetic PnTX G (purity > 97%)	IP Number of animals not specified	65.8		J. Molgó (personal communication)	ToxRTool: 16
PnTX H*	IP Number of animals not specified Fasting not specified	67 CI95: 63-79		Selwood <i>et al.</i> , 2014	ToxRTool: 4 This low score is due to the fact that a lot of details are missing in the article, but this team has already described the protocol for other PnTXs
Portimine*	IP Number of animals not	1570 CI95: 1269-3080	No effect at 500 and 700	Selwood <i>et al.</i> , 2013	ToxRTool: 4 This low score is due to the fact

Toxin (purity)	Route of administration and number of mice	LD50 (µg/kg bw)	MTD (µg/kg bw)	References	Study quality
	specified Fasting not specified				that a lot of details are missing in the article, but this team has already described the protocol for other PnTXs

* Purity verified by NMR according to the authors but percentage not mentioned in the publication

CI95: 95% confidence interval

MTD (Maximum Tolerated Dose): dose at which no mortality or clinical signs are observed

3.2 Subchronic and chronic toxicity

No repeated dose toxicity studies were identified in the literature.

3.3 Mode of action and extrapolation to humans

We reviewed the data available on the mode of action of PnTXs (Delcourt et al., 2019). PnTXs are high-affinity competitive antagonists of nicotinic acetylcholine receptors (nAChRs) (Araoz et al., 2011; Hellyer et al., 2015). Their lethal effects in mice studies are consistent with the inhibition of muscle nAChRs, inducing respiratory distress and paralysis (Delcourt et al., 2019). PnTX E, F, G and A block nerve stimulation-induced skeletal muscle contraction, but do not alter muscle contraction caused by direct stimulation of isolated adult rodent muscles (Hellyer et al., 2013; Benoit et al., 2019).

The clinical effects observed in humans after exposure to compounds (drugs, natural toxins) whose pharmacology is comparable to that of PnTXs have been described (Delcourt et al., 2019). Possible symptoms in humans due to PnTXs could include muscle weakness (myasthenic-like syndrome), dyspnea, anticholinergic syndrome, dysautonomia, pyramidal syndrome, and seizures.

3.4 Development of a health value

The methodology set in ANSES's guide for the development of toxicological reference values (2017) was followed.

3.4.1 Choice of the key study

Two acute oral toxicity studies in mice with purified PnTX G were identified (Munday et al., 2012; ANSES-University of Trieste-CNRS report, 2014 also published in Sosa *et al.*, 2020). The key study selected was the one carried out by the Department of Life Sciences at the University of Trieste with funding from the DGAL and DGS (ANSES-University of Trieste-CNRS report, 2014; Sosa et al., 2020), which obtained the highest score (17 out of 21 with ToxRTool, corresponding to a Klimisch score of 2 - reliable with restriction).

In this study, groups of three to five 4 weeks-old female SD-1 mice received a single administration of purified PnTX G by gavage at a dose of 8, 20, 50, 120, 220, 300, 370 or 450 µg/kg bw. A control group of eight mice was included. Mice were fasted for 3 hours before gavage; food was given back again *ad libitum* 2 hours after administration during the 24-hour observation period. Lethality, clinical signs of toxicity, histological analysis of several organs and blood biochemistry were recorded.

The results regarding lethality and the observed symptoms are shown in Table 2. Administration of PnTX G resulted in mouse lethality from 220 µg PnTX G/kg bw (3/5 mice, in 22 min). No lethality was observed at the doses of 8, 20, 50 and 120 µg/kg bw. All mice (5/5) died at the dose of 370 µg/kg bw (survival time between 13 and 18 min). The LD50 for PnTX G was calculated at 208 µg/kg bw (95% confidence interval = 155-281 µg/kg bw).

Before death, the main symptoms were neurotoxic, i.e. prostration, tremors, jumps, abdominal breathing, hypothermia, hind leg paralysis and cyanosis. No macroscopic organ changes were observed during necropsy of the mice treated with PnTX G. Histological analysis of major organs and tissues revealed only minor changes in the small intestine of mice given PnTX G at doses equal to or greater than 300 µg/kg bw (moderate mucosal degeneration, villous atrophy). No differences in biochemical blood parameters were found between treated and control mice (Sosa *et al.*, 2020).

Table 2: Lethality and clinical signs of toxicity in mice after single administration by gavage of purified PnTX G (ANSES-University of Trieste-CNRS Report, 2014; Sosa *et al.*, 2020)

Purified PnTX G dose	Lethality	Survival time (h:min)	Clinical signs of toxicity
Control	0/8	-	-
8 µg/kg bw	0/3	-	-
20 µg/kg bw	0/3	-	-
50 µg/kg bw	0/3	-	-
120 µg/kg bw	0/3	-	-
220 µg/kg bw	3/5	00:20; 00:22; 00:22	prostration, tremors, jumps, abdominal

			breathing, hypothermia, hind leg paralysis, cyanosis
300 µg/kg bw	4/5	00:12; 00:13; 00:17; 00:23	prostration, tremors, jumps, abdominal breathing, hypothermia, hind leg paralysis, cyanosis
370 µg/kg bw	5/5	00:13; 00:15; 00:16; 00:17; 00:18	prostration, tremors, jumps, abdominal breathing, hypothermia, hind leg paralysis, cyanosis
450 µg/kg bw	5/5	00:12; 00:12; 00:15; 00:16; 00:29	prostration, tremors, jumps, abdominal breathing, hypothermia, hind leg paralysis, cyanosis

3.4.2 Choice of the critical dose

Based on the data from the key study, two approaches were explored:

- First approach: the **120 µg PnTX G/kg bw** dose was selected as the maximum tolerated dose (MTD), i.e. the dose at which no effect is observed in the studied parameters over a 24-hour period after treatment.
- Second approach: calculation of a benchmark dose¹ from a response level of 10% mortality (PROAST Web software, RIVM, v. 65.2). The model that best fitted the experimental data according to the Akaike Information Criterion (AIC) was the probit model, for which the BMDL_{95%} was **69.1 µg PnTX G/kg bw** of mouse.

It is unusual to use data on mouse mortality from single oral administration to derive a health reference value. This is justified by the fact that the mouse mortality occurs rapidly and there is a very small difference between a dose with no signs of toxicity and a lethal dose.

A response level of 10% (for lethality) is considered high but was exceptionally selected in this expert appraisal because the results obtained with a response level of 1% were regarded as too uncertain due to the small number of mice tested in the study.

The results obtained in the other acute oral toxicity study of purified PnTX G with non-fasted mice (Munday *et al.*, 2012) were consistent with those of the key study selected conducted by the Department of Life Sciences at the University of Trieste:

- The LD₅₀ was 150 µg PnTX G/kg bw (95% CI = 105-199 µg/kg bw);

¹ The benchmark dose is a dose producing a measurable effect corresponding to a given response level compared to a control group. The lower limit of its 95% or 90% confidence interval (BMDL_{95%} or BMDL_{90%}) is most often used. This approach is based on modelling of the experimental data taking into account the entire dose-response curve (ANSES 2017a).

- ii. The dose tested without lethality and with no apparent sign of neurotoxicity was 75 µg PnTX G/kg bw.

3.4.3 Choice of uncertainty factors

Uncertainty factors were identified for both approaches explored when establishing the critical dose:

- First approach: the dose of 120 µg PnTX G/kg bw was selected as the maximum tolerated dose (MTD). The associated overall uncertainty factor was 900:
 - inter-species uncertainty factor: $UF_A = 10$
 - inter-individual uncertainty factor: $UF_H = 10$
 - other factors: $UF_D = 3$ (insufficient data) and 3 (to take into account the severity and pattern of the dose-response curve)
- Second approach: the BMDL of 69.1 µg PnTX G/kg bw was chosen as the point of departure. The associated overall uncertainty factor was 525:
 - allometric adjustment factor: 7 for the mouse
 - $UF_A = 2.5$ for the toxicodynamic component
 - $UF_H = 10$
 - $UF_D = 3$ (insufficient data).

3.4.4 Proposed health-based guidance value

Due to insufficient data for the hazard characterisation of PnTX G, the guidance value derived is considered as a **provisional acute benchmark value**.

The provisional acute benchmark value is **0.13 µg PnTX G/kg bw**, (Table 3) regardless of the approach used² (MTD or BMDL).

Table 3: Table summarising development of the provisional acute benchmark value for PnTX G

Critical effect Key study	Critical dose for the mouse	UF	Provisional acute benchmark value for humans	Confidence level
------------------------------	--------------------------------	----	--	---------------------

² MTD approach: $(120 \text{ µg PnTX G/kg bw}) / (10 \times 10 \times 3 \times 3) = 120/900 = 0.1333 \text{ µg PnTX G/kg bw}$
BMDL approach: $69.1 \text{ µg PnTX G/kg bw} / (7 \times 2.5 \times 10 \times 3) = 69.1/525 = 0.1316 \text{ µg PnTX G/kg bw}$

First approach	<i>Critical effect:</i> absence of mortality and symptoms in mice 24 hours after a single oral administration by gavage of purified PnTX G	Maximum tolerated dose of 120 µg PnTX G/kg bw	Total factor: 900 UF _A = 10 UF _H = 10 UF _D = 9 (3 for insufficient data and 3 for the severity and pattern of the dose-response curve).	0.13 µg PnTX G/kg bw	Moderate
Second approach	<i>Key study:</i> ANSES-University of Trieste-CNRS report, 2014; Sosa et al., 2020	BMDL of 69.1 µg PnTX G/kg bw	Total factor: 525 Allometric adjustment: 7 for the mouse UF _A = 2.5 UF _H = 10 UF _D = 3 for insufficient data	0.13 µg PnTX G/kg bw	Moderate

278

279 3.4.5 Proposed maximum tolerable concentration of PnTX G in shellfish

280 Based on the provisional acute benchmark value of 0.13 µg PnTX G/kg bw, a default serving size of
281 400 g of shellfish (EFSA 2010b) and a default body weight of 70 kg, the concentration not to be
282 exceeded in shellfish would be **23 µg PnTX G/kg of total meat**.

283

284 3.4.6 Confidence level

285 An overall confidence level **moderate** was assigned to the provisional acute benchmark value based
286 on the following criteria:

287 - Level of confidence in the type and quality of the body of data: **Low**

288 The literature review revealed that there were few data on the acute oral toxicity of purified PnTXs and
289 none on repeated administration. In fact, only two acute oral toxicity studies conducted on a single
290 species (mice) and sex (females, which are considered to be more sensitive than males) have been
291 published. The lack of data in another rodent species (rats) and of knowledge on the toxicokinetics
292 and toxicodynamics of PnTXs is underlined.

293 - Level of confidence in the choice of the critical effect and the mode of action: **Moderate**

The key study is an acute toxicity study with a single oral administration. Its objective was to study mouse mortality and define a median lethal dose (LD50). Signs of neurological toxicity were also observed and are consistent with the known mode of action of PnTXs.

We stress that it is unusual to use data on mouse mortality from single oral administration to derive a health reference value. This is justified by the fact that the mouse mortality occurs rapidly and there is a very small difference between a dose with no signs of toxicity and a lethal dose.

- Level of confidence in the choice of the key study: **Moderate/High**

The key study was analysed using ToxRTool and obtained a total score of 17 (out of 21), corresponding to a score of 2 using the Klimisch method (reliable study with restriction).

- Level of confidence in the choice of the critical dose: **High**

Two approaches were explored for selecting the critical dose from the key study data. The first approach retained a maximum tolerated dose (MTD), at which there is no mortality or signs of toxicity in mice during the 24-hour observation period (which is considered sufficient because PnTXs are fast-acting toxins, with neurotoxic signs in mice occurring within 30 minutes of administration). The second approach modelled the dose-response relationship (lethality) to calculate a BMDL.

This overall confidence level may be reassessed when new acute oral toxicity data become available for PnTX G or other toxins produced by *V. rugosum*.

4. Occurrence of PnTXs in shellfish

4.1 Ingril lagoon

As soon as PnTX G was identified in France (2011), a retrospective analysis of the samples from 2009 to 2012 was conducted by Ifremer, which revealed the kinetics of shellfish contamination as a function of time. The maximum annual concentrations were 261, 1244, 568 and 652 µg/kg of total mussel meat for 2009, 2010, 2011 and 2012 respectively (Table 4). Concentrations in mussels were higher than those observed in clams, when shellfish were sampled simultaneously (Table 5).

To track this phenomenon and maintain monitoring for PnTXs, sampling of Ingril mussels was carried out monthly between 2013 and 2017 by Ifremer. The results show that PnTX G peaks were observed

between June and September, but the maximum values varied according to the year (887 in 2013, 918 in 2014, 1143 in 2015, 600 in 2016 and 640 in 2017, expressed in µg/kg of total meat).

Table 4: Data on concentrations of PnTX G in mussels from Ingril lagoon from 2010 to 2017 (µg/kg total meat)

Year	2010	2011	2012	2013	2014	2015	2016	2017
Number of samples	30	32	24	12	27	2	13	10
Min	45.4	36.6	11.0	49.5	22.6	572	44.0	52.9
Max	1244	568	652	887	918	1143	600	640
Mean	223	227	209	293	247	858	278	294

Table 5: Data on concentrations of PnTX G in mussels and clams from Ingril lagoon from 2010 to 2012 (µg/kg total meat)

	Mussels	Clams
Number of samples	86	20
Min	11.0	13.6
Max	1244	95.3
Mean	217	28.1

4.2 Other Mediterranean lagoons

In addition to the monitoring carried out in the Ingril lagoon, screening for PnTX G in wild mussel samples took place the same week in four other Mediterranean lagoons in 2013 (Table 6). Maximum levels of PnTX G increased in the following order: Parc Leucate (11 µg/kg of total meat) > Thau (15 µg/kg) > Le Prevost (54 µg/kg) > Vic (89 µg/kg) > Ingril (887 µg/kg).

Table 6: Data on concentrations of PnTX G in mussels from several Mediterranean lagoons in 2013 (µg/kg total meat)

	Parc Leucate	Thau	Le Prévost	Vic	Ingril
Number of samples	12	11	12	11	12
Min	0.68	1.4	11.8	12.3	49.5

Max	10.7	14.7	53.9	89.0	887
Mean	3.4	10.0	26.2	48.4	293

4.3 Atlantic and Corsican coasts

In France, the EMERGTOX scheme for monitoring the emergence of marine biotoxins in shellfish was set up by the DGAL to complement national surveillance schemes for regulated toxins (REPHYTOX, the DGAL's surveillance plan). Its purpose is to bring to light any possible hazard associated with the presence in shellfish of known regulated and unregulated lipophilic toxins, either identified in France or that could be introduced into France via ballast water or commercial trade between countries.

Since 2018, this scheme enables the targeted analysis by chemical methods of PnTXs in mussels and oysters from eleven areas located all along the French metropolitan coasts. During this first year of monitoring, three of these areas (Ingril, Le Scoré in Brittany and the Diana lagoon in Corsica) were affected by the presence of PnTXs, mainly PnTX G and to a lesser degree PnTX A (only in Ingril). Of the three affected areas, the Ingril lagoon remains the most heavily contaminated area (with concentrations detected every month, varying from 40 to 2614 µg PnTX G/kg of digestive gland). The levels found at Le Scoré and Diana were low (maximum around 10 µg PnTX G/kg of digestive gland). The concentrations of PnTX A detected at Ingril ranged from 6 to 32 µg/kg of digestive gland.

We believe it is important to point out that the presence of PnTXs in France is not limited to Mediterranean lagoons. Given that global warming is making ecophysiological conditions more favourable to the development of *V. rugosum*, vigilance should be maintained along the Atlantic coast, since the detection of PnTXs in mussels from Brittany points out the presence of the dinoflagellate producer also in these waters.

4.4 Outside France

The literature review concluded that very limited data on PnTX levels in shellfish from Northern and Southern Europe, Canada and New Zealand have been published. Among the eight studies reviewed (MacKenzie et al., 2011; Rundberget, et al., 2011; McCarron et al., 2012 ; McNabb et al., 2012; Garcia-Altares et al., 2014; Rambla Alegre et al., 2018; Lamas et al., 2019; Otero et al., 2019), the highest reported levels of PnTX G in mussels were 115 µg/kg in Norway (Rundberget et al., 2011), 83 µg/kg in Canada (McCarron et al., 2012) and 59 µg/kg in Spain (Garcia-Altares et al., 2014). These levels are lower than the annual maximum levels reported in France in the Ingril lagoon (between 261

and 1244 µg/kg, depending on the year). It should be noted that the study by Rambla-Alegre *et al.* (2018) reported the persistence of PnTX G in samples of canned mussels (up to 12 µg/kg).

Lastly, ANSES sent a questionnaire to the European Food Safety Authority (EFSA) network of focal points on 15 February 2018, to which 16 Member States replied. At the time of the request, none of the responding health agencies had conducted a health risk assessment for PnTXs. Some respondents provided contamination data in shellfish from their country. Most of the data were already published in a scientific journal and had been identified in our literature search. Some unpolished data were also provided but without information about the performance of the analytical method used.

In conclusion, the concentrations of PnTX G measured in mussels from Ingril are the highest reported in the world to date.

5 Dietary exposure

5.1. Food consumption data

A consumption survey on seafood products in France (CONSOMER) was conducted in 2016-2017 as part of a research agreement between ANSES and CREDOC (2015-CRD-25). The aim of the survey was to assess seafood consumption by an adult population (over 18 years of age) living in coastal areas and with access to local sources of supply. The database includes answers from 2481 adults. Only data for the Mediterranean population were used in the present work, including 821 adults (Table 7).

Table 7: Consumption data of mussels and clams from CONSOMER survey (Mediterranean population only), daily portion size.

	Number of consumers	Mean amount in g/day	P95 in g/day
Mussels	582	91.2	200
Clams	140	19.8	40

5.2 Contamination data

Data on concentrations of PnTX G in mussels and clams from several Mediterranean lagoons from 2010 to 2017 were provided to ANSES by Ifremer (Table 8). Three scenarios were selected, depending on the site where the shellfish were sampled:

- **"All lagoons" scenario:** contamination data of all mussels on the one hand and all clams on the other (from all the lagoons studied);
- **"Ingril" scenario:** contamination data ONLY from the Ingril lagoon, for mussels on the one hand and clams on the other;
- **"Except Ingril" scenario:** contamination data from all lagoons (Vic, le Prévost, Parc Leucate and Thau) except "Ingril", for mussels only (no clams sampled in other lagoons).

Table 8: Concentrations of PnTX G in mussels and clams from several Mediterranean lagoons from 2010 to 2017

	Number of samples of mussels	Concentration of PnTX G (µg/kg total meat)	
		Mean	P95
All lagoons	196	196	633
Ingril	150	249	712
Except Ingril*	46	21.7	60.3
	Number of samples of clams	Concentration of PnTX G (µg/kg total meat)	
		Mean	P95
All lagoons	20	28.1	57
Ingril	20	28.1	57
Except Ingril	0	No data	No data

* Number of samples of mussels in other lagoons than Ingril are described in Table 6.

5.3. Method to calculate exposure

From the individual consumption data and the contamination data, exposure was calculated according to the following equation:

$$E_i = \sum_{k=1}^n \frac{C_{i,k} \times L_k}{BW_i}$$

where:

- E_i is the total daily exposure of individual i associated with the consumption of food k (µg/kg of body weight/day);
- $C_{i,k}$ is the consumption of food k by individual i (kg/d) (serving size for acute exposure);

- L_k is the PnTX G contamination of food k ($\mu\text{g/kg}$ of fresh weight) (95th percentile for acute exposure);
- BW_i is the body weight of individual i (kg).

6 Risk characterisation

6.1 Results

A comparison of acute exposure estimates was made with the provisional acute benchmark value of 0.13 μg PnTX G/kg of body weight. A percentage of the population exceeding this value was calculated, firstly in the total population and secondly for consumers of mussels and clams only (Table 9).

Table 9: Acute exposure associated with consumption of mussels and clams (CONSOMER, Mediterranean area) for adults (in μg PnTX G/kg bw)

	Total population (Mediterranean area)			Consumers only (Mediterranean area)			Provisional acute benchmark value (PABV) exceeded	
	n	Ave_pop	P95_pop	n_cons	Ave_cons	P95_cons	% > PABV	% > PABV_cons
All lagoons	821	0.590	1.572	591	0.816	1.733	71	100
Ingril	821	0.663	1.767	591	0.917	1.946	71	100
Except Ingril (mussels only*)	821	0.056	0.149	582	0.079	0.167	7.3	10.23

* analyses of clams were only carried out for the Ingril lagoon

n: number of adult individuals in the population monitored during the survey (this population includes consumers and non-consumers of mussels and clams from the first column)

Ave_pop: average exposure in the population (whether consumers or not) ($\mu\text{g/kg}$ bw)

P95_pop: exposure to the 95th percentile calculated in the population ($\mu\text{g/kg}$ bw/d)

n_cons: number of adult individuals consuming mussels and clams from the first column (consumers only)

Ave_cons: average exposure in the consumer population only ($\mu\text{g/kg}$ bw/day)

P95_cons: exposure to the 95th percentile calculated in the consumer population only ($\mu\text{g/kg}$ bw/day)

% > PABV: percentage of the general population exceeding the provisional acute benchmark value of 0.13 $\mu\text{g/kg}$ bw

% > PABV_cons: percentage of consumer population only exceeding the provisional acute benchmark value of 0.13 $\mu\text{g/kg}$ bw

These results show that in the tested scenarios including the contamination data from the Ingril lagoon the provisional acute benchmark value would be exceeded in 71% of adults according to the consumption data from the CONSUMER study for the Mediterranean area (in the scenario including non-consumers of mussels and clams). If only individuals consuming mussels and clams are considered, this proportion increases to 100%. In the tested scenario excluding the contamination data from the Ingril lagoon, the provisional acute benchmark value would be exceeded in 7% of the total Mediterranean population and in 10% of the population of consumers.

6.2 Conclusion in terms of health concern

Considering the percentage of the French population for which the provisional acute benchmark value would be exceeded in the "Except Ingril" scenario, there may be a health concern related to the consumption of shellfish contaminated with PnTXs from these Mediterranean lagoons. Estimates were not made for the other French sites where PnTXs were detected in mussels (Le Scoré, Diana) because the contamination was measured only in the digestive glands, and the distribution of PnTXs between the digestive gland and the total meat is not known.

7. Recommendations for surveillance

7.1 Environmental surveillance

Several studies on the ecology of the dinoflagellate *V. rugosum* have been carried out *in vitro* and *in situ*. However, its origin and the determinism of its blooms in natural environments remain to be elucidated. While the contamination of molluscs by PnTXs in some areas is undeniable, it remains difficult to establish the relationship with *V. rugosum* blooms as the observation of the pelagic phase in the water column is challenging. Official surveillance of shellfish production areas based on the identification and counting of toxic phytoplanktonic species in the water column is therefore ill-suited to this species.

Several monitoring options can be proposed:

- 1) Regarding dinoflagellate surveillance, monitoring of the benthic population of this organism in risk areas is recommended. It is essential to sample the macroalgae present in the area, in order to collect *V. rugosum* cells. The protocol set up for "*Ostreopsis*" monitoring and tested in the study by Abadie *et al.* (2018) shows that such methodology is adequate.

2) Alongside the monitoring of *V. rugosum*, more systematic surveillance of the presence of PnTX A and G toxins in molluscs should be considered. The extraction protocol for PnTXs is identical to the one used for official monitoring of lipophilic toxins in the French shellfish production areas (REPHYTOX). Although an additional step for LC-MS/MS analysis is required, the cost remains moderate. A systematic screening for PnTXs in total meat could be added to the REPHYTOX analyses of lipophilic toxins, and not only on samples taken for the EMERGTOX scheme, that focuses only on the digestive gland.

Such proposal will enable estimating and monitoring the presence of PnTXs in bivalve molluscs in national production areas.

7.2 Health monitoring, reporting and procedures

To carry on with this issue, ANSES is working on raising awareness among health professionals of the potential neurological symptoms that could be associated with PnTXs poisoning in humans (Delcourt et al., 2019). An epidemiological study is currently underway with the objective to identify potential human cases related to shellfish consumption and PnTX exposure in France.

8. Conclusion

This work points out that, in case of high consumption and/or high contamination, the provisional acute benchmark value for PnTX G could be exceeded, suggesting a health concern related to the consumption of shellfish contaminated with PnTXs from Mediterranean lagoons, and particularly from the Ingril area. There is currently no shellfish production intended for sale from Ingril area. Nevertheless, ANSES highlighted the need to avoid all consumption of shellfish from this area. ANSES recommends that PnTXs should be taken into account in the French official monitoring of shellfish production areas.

The risk assessment underlying this conclusion is based on a worst-case scenario for estimating exposure to PnTX G, using the 95th percentile of shellfish contamination. A more realistic risk assessment could be conducted using a probabilistic approach (taking into account the distribution of all the data, on both contamination and consumption), which would require the acquisition of more contamination data in shellfish, particularly oysters (Anses, 2019).

9. Data gaps

ANSES recommends undertaking research to estimate more accurately the exposure of shellfish consumers. More data on shellfish contamination by PnTXs are needed, including other species than mussels and clams. More research is also needed to better characterize the oral toxicity of toxins produced by *V. rugosum*. As portimine is produced in large quantities by Mediterranean strains of *Vulcanodinium rugosum*, this toxin should be also considered. This toxin only seems to accumulate at very low levels in shellfish. Additional data are required to confirm its low accumulation in shellfish. Although less toxic than PnTXs in mice by the intraperitoneal route, the toxicological data for portimine are limited and investigation should be carried on. Three of the identified PtTXs (A to C) are structural analogues belonging to the PnTX group but, due to the lack of data available in the literature, they could not be taken into account in this risk assessment. More toxicity and contamination studies are needed.

An extensive acute oral toxicity study on PnTX G in rodents is needed (including description of clinical symptoms, assessment of biological, hematological and anatomopathological parameters as in the repeated dose studies, for a range of doses close to the maximum tolerated dose in the 2014 ANSES-University of Trieste-CNRS report study/Sosa et al., 2020) , with a 14-day observation period.

A 28-day repeated oral administration study (OECD guideline 407) or even a 90-day study (OECD guideline 408) should be conducted. Given the data currently available, the effects on the central and peripheral nervous system as well as on the cardiovascular system should be investigated. Conducting a neurotoxicity study according to OECD guideline 424 is recommended. Preliminary results on *in ovo* exposure of chickens (Couesnon et al., 2014; unpublished results) suggest that PnTXs could affect embryonal growth. Therefore, the effects of PnTXs on offspring should be investigated, according to the OECD guideline 422.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of Interest: The authors declare no conflict of interest. The authors' declarations of interests are made public via the ANSES website (www.anses.fr).

References

Abadie, E., A. Muguet, T. Berteaux, N. Chomerat, P. Hess, E. Roque D'OrbCastel, E. Masseret, M. Laabir, 2016. Toxin and Growth Responses of the Neurotoxic Dinoflagellate *Vulcanodinium rugosum* to Varying Temperature and Salinity. *Toxins* 8(5). doi: 10.3390/toxins8050136.

Abadie, E., C. Chiantella, A. Crottier, L. Rhodes, E. Masseret, T. Berteaux, M. Laabir, 2018. What are the main environmental factors driving the development of the neurotoxic dinoflagellate *Vulcanodinium rugosum* in a Mediterranean ecosystem (Ingril lagoon, France)? *Harmful Algae* 75:75-86. doi:10.1016/j.hal.2018.03.012.

Al Muftah, A., A.I. Selwood, A.J. Foss, H.M. Al-Jabri, M. Potts, M. Yilmaz, 2016. Algal toxins and producers in the marine waters of Qatar, Arabian Gulf. *Toxicon* 122:54-66.

Anses (Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail), 2017.

Valeurs toxicologiques de référence. Guide d'élaboration de l'Anses (only in French).

<https://www.anses.fr/fr/system/files/SUBSTANCES2017SA0016Ra.pdf>

Anses (Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail), 2019. Risques liés aux pinnatoxines dans les coquillages. Avis de l'Anses et Rapport d'expertise collective, Mars 2019 - Édition scientifique, 136 p, <https://www.anses.fr/fr/system/files/ERCA2016SA0013Ra.pdf> (full report only in French).

Anses-Université de Trieste-CNRS, 2014. Pinnatoxines en lien avec l'espèce *Vulcanodinium rugosum* (II) Partie 2 : Etudes sur la toxicité des pinnatoxines et d'extraits de *Vulcanodinium rugosum*, 72p. <https://archimer.ifremer.fr/doc/00285/39635/38127.pdf>

559 Araoz, R., D. Servent, J. Molgó, B.I. Iorga, C. Fruchart-Gaillard, E. Benoit, Z. Gu, C. Stivala. A.
 560 Zakarian. 2011, Total synthesis of pinnatoxins A and G and revision of the mode of action of
 561 pinnatoxin A. J. Am. Chem. Soc. 133:10499-10511.
 562
 563 Benoit, E., A. Couesnon, J. Lindovsky, B.I. Iorga, R. Araoz, D. Servent, .A. Zakarian, J. Molgó. 2019.
 564 Synthetic pinnatoxins A and G reversibly block mouse skeletal neuromuscular transmission In Vivo
 565 and In Vitro. Mar. Drugs 17(5). pii: E306. doi: 10.3390/md17050306.
 566 Couesnon, A., J. Lindovsky, A. Zakarian, S. Creuzet, J. Molgó, 2014. Pinnatoxins block skeletal
 567 neuromuscular junction activity and affect embryo development. Toxicon 91:175-176.
 568 doi:10.1016/j.toxicon.2014.08.036.
 569
 570 Chou, T., T. Haino, M. Kuramoto, D. Uemura, 1996. Isolation and structure of pinnatoxin D, a new
 571 shellfish poison from the Okinawan bivalve *Pinna muricata*. Tetrahedron Letters 37(23): 4027-4030.
 572
 573 Delcourt, N., Lagrange, E., Abadie, E., Fessard, V., Frémy, J.-M., Vernoux, J.-P., Peyrat, M.-B.,
 574 Maignien, T., Arnich, N., Molgó, J., Mattei, C., 2019. Pinnatoxins' deleterious effects on cholinergic
 575 networks: From experimental models to human health. Marine Drugs 17, Issue 7, Article number 425.
 576
 577 EFSA (European Food Safety Authority). 2010a. Scientific Opinion on marine biotoxins in shellfish –
 578 Cyclic imines (spirolides, gymnodimines, pinnatoxins and pteriattoxins). EFSA Journal 8 (6):1628. doi:
 579 doi:10.2903/j.efsa.2010.1628.
 580
 581 EFSA (European Food Safety Authority). 2010b. Statement on further elaboration of the consumption
 582 figure of 400 g shellfish meat on the basis of new consumption data. EFSA Journal 8 (8):1706.
 583 doi:10.2903/j.efsa.2010.1706.
 584
 585 Garcia-Altares, M., A. Casanova, V. Bane, J. Diogene, A. Furey, P. de la Iglesia, 2014. Confirmation of
 586 pinnatoxins and spirolides in shellfish and passive samplers from Catalonia (Spain) by liquid
 587 chromatography coupled with triple quadrupole and high-resolution hybrid tandem mass spectrometry.
 588 Mar Drugs 12 (6):3706-32. doi: 10.3390/md12063706.

589

590 Garrett, M. J., C. Puchulutegui, A. I. Selwood, J. L. Wolny. 2014. Identification of the harmful
 591 dinoflagellate *Vulcanodinium rugosum* recovered from a ballast tank of a globally traveled ship in Port
 592 Tampa Bay, Florida, USA." *Harmful Algae* 39:202-209. doi: 10.1016/j.hal.2014.07.014.

593

594 Hellyer, S. D., A. I. Selwood, L. Rhodes, D. S. Kerr, 2013. Neuromuscular blocking activity of
 595 pinnatoxins
 596 E, F and G. *Toxicon* 76:214-20. doi: 10.1016/j.toxicon.2013.10.009.

597

598 Hellyer, S.D., D. Indurthi, T. Balle, V. Runder-Varga, A.I. Selwood, J.D.Tyndall, M. Chebib, L. Rhodes,
 599 D.S. Kerr. 2015. Pinnatoxins E, F and G target multiple nicotinic receptor subtypes. *J. Neurochem.*
 600 135:479-91. doi: 10.1111/jnc.13245.

601

602 Hernandez-Becerril, D. U., M. C. Rodriguez-Palacio, C. Lozano-Ramirez, 2013. Morphology and life
 603 stages of the potentially pinnatoxin-producing thecate dinoflagellate *Vulcanodinium rugosum* from the
 604 tropical Mexican Pacific. *Botanica Marina* 56 (5-6):535-540. doi: 10.1515/bot-2013-0079.

605

606 Ifremer (Institut français de recherche pour l'exploitation de la mer). 2012. P. Hess, F. Hervé, E.
 607 Abadie, V. Séchet, J. Molgó, Z. Amzil, V. Fessard. Pinnatoxines en lien avec l'espèce *Vulcanodinium*
 608 *rugosum*. Référence Ifremer: 11/1219401/NYF, 09-2012 - R.RBE/EMP/PHYC 12-05, 144 p.
 609 <https://archimer.ifremer.fr/doc/00094/20518/18190.pdf>

610

611 Klimisch, H. J., M. Andreae, U. Tillmann, 1997. A systematic approach for evaluating the quality of
 612 experimental toxicological and ecotoxicological data." *Regul Toxicol Pharmacol* 25 (1):1-5.
 613 doi:10.1006/rtph.1996.1076.

614

615 Lamas, J.P., Arévalo, F., Moroño, Á., Correa, J., Muñíz, S., Blanco, J. 2019. Detection and spatio-
 616 temporal distribution of pinnatoxins in shellfish from the Atlantic and Cantabrian coasts of Spain.
 617 *Toxins* 11(6). pii: E340. doi: 10.3390/toxins11060340.

618

619 MacKenzie, L. A., A. I. Selwood, P. McNabb, L. Rhodes, 2011. Benthic dinoflagellate toxins in two
620 warmtemperate estuaries: Rangaunu and Parengarenga Harbours, Northland, New Zealand. *Harmful*
621 *Algae* 10 (6):559-566. doi: 10.1016/j.hal.2011.02.007.

622

623 McCarron, P., W. A. Rourke, W. Hardstaff, B. Pooley, M. A. Quilliam, 2012. Identification of
624 pinnatoxins
625 and discovery of their fatty acid ester metabolites in mussels (*Mytilus edulis*) from eastern Canada. *J*
626 *Agric Food Chem* 60 (6):1437-46. doi: 10.1021/jf204824s.

627

628 McCauley, J.A., K. Nagasawa, P.A., S.G. Mischke, M.A. Semones, Y. Kishi, 1998. Total synthesis of
629 pinnatoxin A. *J. Am. Chem. Soc.* 120(30):7647-7648.

630

631 McNabb, P. S., D. J. McCoubrey, L. Rhodes, K. Smith, A. I. Selwood, R. van Ginkel, A. L. MacKenzie,
632 R. Munday, P. T. Holland, 2012. New perspectives on biotoxin detection in Rangaunu Harbour, New
633 Zealand arising from the discovery of pinnatoxins. *Harmful Algae* 13:34-39.
634 doi:10.1016/j.hal.2011.09.013.

635

636 Molgó, J., P. Marchot, R. Aráoz, E. Benoit, B.I. Iorga, A. Zakarian, P. Taylor, Y. Bourne, D. Servent.
637 2017. Cyclic imine toxins from dinoflagellates: a growing family of potent antagonists of the nicotinic
638 acetylcholine receptors. *J. Neurochem.*, 142 S2:41-51. doi: 10.1111/jnc.13995.

639

640 Moreira-Gonzalez, A.R., A. Comas-Gonzalez, A. Valle-Pombrol, M. Seisdedo-Losa, C. Alonso-
641 hernandez, O. Hernandez-Leyva, L.F. Ferandes, N. Chomerat, G. Bilien, F. Herve, G.A. Rovillon, P.
642 Hess, L.L. Mafra, 2018. Summer bloom of *Vulcanodinium rugosum* in Cienfuegos Bay (Cuba)
643 associated to dermatitis in swimmers. 18th edition of the International Conference on Harmful Algae
644 (ICHA), 21- 26 October 2018, Nantes (France). Poster 238. Abstract book
645 https://www.icha2018.com/medias/content/files/ICHA_2018_Abstract_Book_v8.pdf

646

647 Munday, R., A. I. Selwood, L. Rhodes, 2012. Acute toxicity of pinnatoxins E, F and G to mice. *Toxicon*
648 60 (6):995-999.

649

650 Nézan, E., Chomérat, N., 2011. *Vulcanodinium rugosum* gen. nov., sp. nov. (dinophyceae): A new
 651 marine dinoflagellate from the French Mediterranean coast. *Cryptogam. Algal.* 32, 3–18.

652

653 Otero, P., Miguéns, N., Rodríguez, I., Botana, L.M. 2019. LC-MS/MS Analysis of the Emerging Toxin
 654 Pinnatoxin-G and High Levels of Esterified OA Group Toxins in Galician Commercial Mussels. *Toxins*
 655 11(7), 394. doi: 10.3390/toxins11070394.

656

657 Rambla-Alegre, M., C. O. Miles, P. de la Iglesia, M. Fernandez-Tejedor, S. Jacobs, I. Sioen, W.
 658 Verbeke, I. A. Samdal, M. Sandvik, V. Barbosa, A. Tediosi, E. Madorran, K. Granby, M. Kotterman, T.
 659 Calis, J. Diogene, 2018. Occurrence of cyclic imines in European commercial seafood and consumers
 660 risk
 661 assessment. *Environ Res* 161:392-398. doi: 10.1016/j.envres.2017.11.028.

662

663 Rhodes, L., K. Smith, A. Selwood, P. McNabb, R. van Ginkel, P. Holland, R. Munday, 2010.
 664 Production of pinnatoxins by a peridinoid dinoflagellate isolated from Northland, New
 665 Zealand. *Harmful Algae* 9(4):384-389. doi: 10.1016/j.hal.2010.01.008.

666

667 Rhodes, L., K. Smith, A. Selwood, P. McNabb, S. Molenaar, R. Munday, C. Wilkinson, G. Hallegraeff,
 668 2011. Production of pinnatoxins E, F and G by scrippsielloid dinoflagellates isolated from Franklin
 669 Harbour, South Australia. *New Zealand Journal of Marine and Freshwater Research* 45 (4):703-709.
 670 doi: 10.1080/00288330.2011.586041.

671

672 Rundberget, T., J. A. Aasen, A. I. Selwood, C. O. Miles, 2011. Pinnatoxins and spirolides in Norwegian
 673 blue mussels and seawater. *Toxicon* 58 (8):700-11. doi: 10.1016/j.toxicon.2011.08.008.

674

675 Selwood, A. I., C. O. Miles, A. L. Wilkins, R. van Ginkel, R. Munday, F. Rise, P. McNabb, 2010.
 676 Isolation, structural determination and acute toxicity of pinnatoxins E, F and G. *J Agric Food Chem*
 677 58(10):6532-42. doi: 10.1021/jf100267a.

678

679 Selwood, A. I., A. L. Wilkins, R. Munday, F. Shi, L. L. Rhodes, P. T. Holland, 2013. Portimine: A
680 bioactive metabolite from the benthic dinoflagellate *Vulcanodinium rugosum*. *Tetrahedron Letters*
681 54(35):4705-4707. doi: 10.1016/j.tetlet.2013.06.098.

682

683 Selwood, A. I., A. L. Wilkins, R. Munday, H. Gu, K. F. Smith, L. L. Rhodes, F. Rise, 2014. Pinnatoxin
684 H:
685 A new pinnatoxin analogue from a South China Sea *Vulcanodinium rugosum* isolate. *Tetrahedron*
686 *Letters* 55 (40):5508-5510. doi: 10.1016/j.tetlet.2014.08.056.

687

688 Smith, K. F., L. L. Rhodes, S. Suda, A. I. Selwood, 2011. A dinoflagellate producer of pinnatoxin G,
689 isolated from sub-tropical Japanese waters." *Harmful Algae* 10 (6):702-705.
690 doi:10.1016/j.hal.2011.05.006.

691

692 Sosa, S., Pelin, M., Cavion, F., Hervé, F., Hess, P., Tubaro, A. 2020. Acute oral toxicity of pinnatoxin
693 G in mice. *Toxins (Basel)*. 2020 Jan 28;12(2). pii: E87. doi: 10.3390/toxins12020087.

694

695 Takada, N., N. Umemura, K. Suenaga, D. Uemura, 2001. Structural determination of pteriatoxins A, B
696 and C, extremely potent toxins from the bivalve *Pteria penguin*. *Tetrahedron Letters* 42 (20):3495-
697 3497. doi: 10.1016/S0040-4039(01)00478-6.

698

699 Uemura, D., T. Chou, T. Haino, A. Nagatsu, S. Fukuzawa, S. Z. Zheng, H. S. Chen, 1995. Pinnatoxin-
700 a -a toxic amphoteric macrocycle from the Okinawan bivalve *Pinna muricata*. *J Am Chem Soc*
701 117(3):1155-1156. doi: 10.1021/ja00108a043.

702

703 Zendong, Z., C. Herrenknecht, E. Abadie, C. Brissard, C. Tixier, F. Mondegue, V. Sechet, Z. Amzil, P.
704 Hess. 2014. Extended evaluation of polymeric and lipophilic sorbents for passive sampling of marine
705 toxins. *Toxicon* 91:57-68. doi: 10.1016/j.toxicon.2014.03.010.

706

707 Zeng, N., H. Gu, K. F. Smith, L. L. Rhodes, A. I. Selwood, W. Yang, 2012. The first report of
708 *Vulcanodinium rugosum* (Dinophyceae) from the South China Sea with a focus on the life cycle. *New*

709 Zealand Journal of Marine and Freshwater Research 46 (4):511-521.
710 doi:10.1080/00288330.2012.719911.