



Detection of norovirus, hepatitis A and hepatitis E viruses in multicomponent foodstuffs

Catherine Hennechart-Collette, Océane Dehan, Michel Laurentie, Audrey Fraisse, Sandra Martin-Latil, Sylvie Perelle

► To cite this version:

Catherine Hennechart-Collette, Océane Dehan, Michel Laurentie, Audrey Fraisse, Sandra Martin-Latil, et al.. Detection of norovirus, hepatitis A and hepatitis E viruses in multicomponent foodstuffs. International Journal of Food Microbiology, 2021, 337, pp.108931. 10.1016/j.ijfoodmicro.2020.108931 . anses-03015846

HAL Id: anses-03015846

<https://anses.hal.science/anses-03015846>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Detection of norovirus, hepatitis A and hepatitis E viruses in multicomponent foodstuffs

**Catherine Hennechart-Collette¹, Océane Dehan¹, Michel Laurentie², Audrey
Fraisie¹, Sandra Martin-Latil¹, Sylvie Perelle^{1*}**

**¹Université Paris-Est, ANSES Laboratory for Food Safety, F-94700 Maisons-Alfort,
France. ²Anses Fougères laboratory, F-35306 Fougères cedex**

*** Corresponding author. Tel: +33 (0)1 49 77 27 99; fax: +33 (0)1 43 68 97 62.**

***E-mail address:* sylvie.perelle@anses.fr**

Abstract

Among the enteric viruses implicated in foodborne outbreaks, the human norovirus and hepatitis viruses A and E (HAV and HEV) represent a serious public health concern. International standard ISO 15216 proposes methods for detecting HAV and norovirus (genogroups I and II) RNA from soft fruit, leaf, stem and bulb vegetables, bottled water or food surfaces. These methods had not previously been validated for detecting the targeted viruses in other foodstuffs such as multicomponent foods, nor for detecting other viruses in foodstuffs. The aim of this study was to characterise a method derived from the vegetable method described in ISO 15216 to detect HAV, HEV and norovirus in artificially-contaminated multicomponent foodstuffs according to the recent international standard ISO 16140-4.

Results showed that the mean recovery rates for all settings did not differ according to the operator. The mean extraction yields ranged from 0.35% to 40.44% for HAV, 5.19% to 100% for HEV, 0.10% to 40.61% for norovirus GI and 0.88% to 69.16% for norovirus GII. The LOD₉₅ was 10² genome copies/g for HAV, HEV and norovirus GII and 10³ genome copies/g for norovirus GI. The LOQ was 2.90×10⁴, 1.40×10³, 1.60×10⁴ and 1.30×10⁴ genome copies/g for HAV, HEV, norovirus GI and norovirus GII respectively. The MNV-1 process control was detected in 120 out of 128 RNA extracts analysed and was recovered with an efficiency of between 3.83% and 50.22%. The mean inhibition rates of quantitative real-time RT-PCR reaction ranged from 3.25% to 28.70% and varied significantly with the type of food matrix. The described method could be used to detect viruses in composite food products for routine diagnosis needs.

Keywords: Multicomponent foodstuff; Human norovirus; Hepatitis virus (A, E); quantitative real-time RT-PCR; Detection; Process control

1. Introduction

Viruses are a leading cause of foodborne disease worldwide. Human norovirus and hepatitis viruses (hepatitis A (HAV) and hepatitis E (HEV)) are recognised to be the main viruses of public health importance. Enteric viruses are primarily transmitted via the faecal-oral and vomit-oral routes, including direct person-to-person contact, consumption of contaminated food or water, contact with contaminated environmental surfaces (Kotwal and Cannon, 2014; Matthews et al., 2012) and for hepatitis E virus direct contact with infected animals (Dalton et al., 2013; Meng, 2010; Pavio et al., 2006, 2010). Various food products, such as bivalve molluscs, fresh fruit and vegetables including different types of lettuce, onions or berries have been involved in foodborne disease outbreaks worldwide (Bernard et al., 2014; Donnan et al., 2012; Ethelberg et al., 2010; Fournet et al., 2012; Gallot et al., 2011; Herman et al., 2015; Le Guyader et al., 2010; Muller et al., 2016; Sarvikivi et al., 2012; Thebault et al., 2013; Wadl et al., 2010). A wide variety of foodstuffs have been implicated and mixed foods and ready to eat meals have frequently been considered as the carriers. In France, in addition to the high-risk food categories, meat, water, dishes with mixed foods or ready to eat meals have also been found to be responsible for 33% of foodborne illnesses and more specifically for 31% of viral foodborne disease outbreaks (*Santé publique France*, 2017). Virus outbreaks have been much more frequently reported in settings using catering services and in restaurants (*Santé publique France*, 2017). Food contamination mainly occurs in restaurants during food preparation by infected workers, and is most often associated with

food service settings (Baert et al., 2008; Barrabeig et al., 2010; Franck et al., 2015; Ronnqvist et al., 2014; Stals et al., 2013).

Most foodborne viruses are currently difficult or impossible to cultivate (Hamza et al., 2011), and sensitive molecular methods are therefore used to detect them in food. International standard ISO 15216 proposes methods for detecting HAV and norovirus (genogroup I and II) RNA from soft fruit, leaf, stem and bulb vegetables, bottled water or food surfaces. However, these methods have neither been validated to detect **other targeted viruses in other foodstuffs, such as multicomponent foods.**

The aim of this study was to validate a method adapted from ISO 15216-1 for the detection of HAV, HEV and norovirus in multicomponent foodstuffs based on the recent international standard ISO 16140-4 (Microbiology of the food chain — Method validation —Part 4: Protocol for method validation in a single laboratory) (Anonymous, 2017) to ensure the safety of these products.

2. Materials and methods

2.1. Viruses and cells

The FRhK-4 (foetal rhesus monkey kidney) cell line was purchased from the American Type Culture Collection (ATCC) (ATCC® CRL-1688™) (LGC standards SARL, Illkirch, France). These epithelial cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco™) supplemented with non-essential amino acids (NEAA, Gibco™) and 10% of heat-inactivated foetal bovine serum (FBS, Gibco™) (Thermo Fisher Scientific, Waltham, MA, 209 USA). Cells were maintained at 37°C in a humidified atmosphere containing 95% air and 5% CO₂.

HAV strain HM175/18f, clone B (VR-1402), was obtained from ATCC. This clone replicates rapidly and has cytopathic effects in cell cultures (Lemon et al., 1991). HAV stock was produced by propagation in foetal rhesus monkey kidney (FRhK-4) cells (ATCC, CRL-1688) (Cromeans et al., 1987). The virus production titre was determined in HAV RNA genomic copies using a **quantitative real-time RT-PCR** standard curve obtained from the ten-fold diluted *in vitro* RNA transcripts as previously described (Fraisse et al., 2017). The HAV stock had a titre of 2.90×10^9 genome copies/mL.

A clarified HEV genotype 3e suspension was obtained from faecal samples of infected swine provided by the ANSES Maisons-Alfort Laboratory for Animal Health. The partial sequence of ORF2 had previously been deposited with GenBank (accession number JF718793). The faecal sample was suspended in 10 mM Phosphate Buffered Saline (PBS), pH 7.4, to obtain a final 10% suspension (w/v), vortexed and centrifuged at 4000 g for 20 min at 4°C. Aliquots of supernatants containing viral particles were then stored at -80°C. The number of HEV RNA copies in the faecal suspension was quantified by using the standard curve obtained with the quantified *in vitro* RNA transcripts as previously described (Martin-Latil et al., 2012). The clarified suspension stock of HEV had a titre of 1.40×10^7 genome copies/mL.

Stool samples of norovirus GI (E8050) and norovirus GII (E7022) from infected humans were provided by the French National Reference Center for Gastroenteritis Viruses in Dijon, France. The faecal samples were suspended in 10 mM **phosphate-buffered saline** (PBS), pH 7.4 to obtain a final 10% suspension (w/v), and then vortexed and centrifuged at 4,000 x g for 20 min at 4°C. Aliquots of 100 µL were frozen and kept at -80°C for later use. The genomic titres of the clarified faecal suspensions were determined by **quantitative real-time**

RT-PCR using a quantitative real-time RT-PCR standard curve obtained with the 10-fold diluted *in vitro* RNA transcripts as previously described (Hennechart-Collette et al., 2014). The clarified suspension stocks of norovirus GI and norovirus GII had titres of approximately 1.6×10^7 and 1.3×10^7 genome copies/mL respectively.

The murine norovirus MNV-1 (CW1 strain) was provided to the ANSES Fougères Laboratory (Fougères, France) by Dr. H. Virgin from Washington University (Saint Louis, MO, USA), and was propagated in a mouse leukaemic monocyte macrophage (RAW 264.7, ATCC TIB-71) cell line (Cannon et al., 2006). RAW 264.7 was grown at 37°C in an atmosphere containing 5% CO₂ in DMEM supplemented with GlutaMAX™, 1% non-essential amino acids and 10% foetal bovine serum (Life Technologies, Saint Aubin, France). The production stock of MNV-1 had titres of approximately 2.15×10^7 TCID₅₀/mL.

2.2. Experimental design and artificial contamination of multicomponent foodstuffs

Twelve multicomponent foodstuff samples (samples 1 to 12 including vegetables, mixed vegetables, meals including meat or fish, soup or sauce) were purchased from a local supermarket. These samples were selected by taking account of food processing factors such as freezing, vacuum packaging or canning. Table 1 describes the selected samples according to matrix type and processing factor. The selected food samples were classified according to the EFSA FoodEx2 classification (https://data.food.gov.uk/codes/foodtype/hierarchy/main/_report).

The experimental design described in ISO 16140-4:2017 was used. Multicomponent foods were artificially contaminated at four contamination levels for all four viruses (HAV, HEV,

norovirus GI and GII). The concentration levels for HAV, HEV and norovirus were obtained by using different inoculum dilution levels. The four inoculation levels ranged from 2.9×10^3 to 2.9×10^6 genome copies for HAV, 1.4×10^2 to 1.4×10^5 genome copies for HEV, 1.6×10^3 to 1.6×10^6 genome copies for norovirus GI and 1.3×10^3 to 1.3×10^6 genome copies for norovirus GII. The food samples were randomly allocated to four different settings (R1, R2, R3 and R4). For each setting, four different matrices per contamination level were analysed by two operators (operators A and B). Table 2 describes the allocation of the food matrices to the different settings along with virus level. Thus, for each virus, four settings (R1 to R4) with four inoculation levels and two replicates (operators A and B) were tested corresponding to 32 analyses in total. Tests were conducted on three different days to evaluate the reproducibility of the method. R1 and R2 settings were analysed on the same day, R3 and R4 on two different days.

Each sample was co-inoculated with 2.15×10^4 TCID₅₀ of MNV-1 (process control virus) just before adding elution buffer. For each matrix, one food sample inoculated only with sterile water was used as a negative control during the entire sample processing and viral detection procedure.

2.3. Sample processing

The method used to recover the viruses was adapted from the ISO 15216-1 procedure described for vegetables by adding a final purification step with chloroform-butanol. Briefly, each inoculated sample (25 g) placed in a 400 mL polypropylene bag containing a filter compartment was soaked in 40 mL of elution buffer (Tris-HCl 100 mM, glycine 50 mM, 1% beef extract (Sigma-Aldrich, Saint-Quentin Fallavier, France)) at pH of 9.5. The rinsing fluid was removed via the bag's filter compartment and centrifuged at 10,000 g for 30 min at 4°C

to pellet the food particles. The pH of the decanted supernatant was adjusted to 7.2 ± 0.2 by the addition of 5 N HCl while the fluid was swirled constantly. The neutralised supernatant was supplemented with 10% (w/v) polyethylene glycol (PEG) 8,000 and 0.3 M NaCl (Sigma-Aldrich), and was then incubated **with constant shaking** at 4°C for one hour. The viruses were concentrated by centrifuging the solution at 10,000 g for 30 min at 4°C. The supernatant was discarded then what remained was centrifuged again at 10,000 g for 5 min at 4°C to compact the pellet. The pellet was then suspended in 500 µL of PBS and vortexed with 500 µL of chloroform: butanol, 1:1 (v/v). The suspension was then incubated for 5 min at room temperature, and centrifuged at 8,000 g for 15 min at 4°C. The upper aqueous phase containing viruses was directly processed by the nucleic acid extraction procedure.

2.4. Viral RNA extraction

NucliSENS® easyMAG™ lysis buffer (BioMérieux, Marcy l'Etoile, France) was added to the virus suspension (up to 3 mL) and total nucleic acid was extracted using the NucliSENS® easyMAG™ platform with the “off-board Specific A” protocol according to the manufacturer’s instructions. Nucleic acids were eluted in 100 µL of elution buffer and stored at -80°C.

2.5. Primers and probes

The primers and probes used to quantify HAV, norovirus GI and norovirus GII have already been described in the literature (Costafreda et al., 2006; da Silva et al., 2007; Kageyama et al., 2003; Loisy et al., 2005; Pinto et al., 2009; Svraka et al., 2007) and are recommended in

ISO 15216 standards. The primers and probe used to quantify HEV were adapted from the model described by Jothikumar et al. (2006) and have been previously described (Martin-Latil et al., 2012b, 2014). The primers and the TaqMan® probe targeting the ORF1 polyprotein of the murine norovirus (MNV-1), which were designed using Beacon Designer software (Bio-Rad, Marnes-la-Coquette, France), have also been described in the literature (Martin-Latil et al., 2012b). All the primers and probes were purchased from Eurofins MWG Operon (Les Ulis, France).

2.6. Quantitative real-time RT-PCR conditions

One-step quantitative real-time RT-PCR amplifications were performed in duplicate on the CFX96™ real-time PCR detection system (Bio-Rad). Reactions were performed in a 25 µL reaction mixture containing 1X of RNA UltraSense™ master mix and 1.25 µL of RNA UltraSense™ enzyme mix, which are components of the RNA UltraSense™ One-Step Quantitative RT-PCR System (Life Technologies), 2 U RNase inhibitor (Life Technologies), 1.25µg of bovine serum albumin (Life Technologies), 500 nM of forward primer, 900 nM of reverse primer, 250 nM of probe and 5 µL of RNA extract. Positive controls containing RNA extracted from virus suspensions and a negative control containing all the reagents except the RNA template were included with each set of reaction mixtures. The one-step quantitative real-time RT-PCR programme involved 60 min of reverse transcription of RNA at 55°C, followed by a 5 min denaturation step at 95°C, and finally 45 cycles of 15 s at 95°C, 1 min at 60°C and 1 min at 65°C. Fluorescence was recorded by the apparatus at the end of the elongation steps (1 min at 65°C) for each amplification cycle. All the samples were characterised by a corresponding cycle threshold (Ct) values. Negative samples gave no Ct

value. A standard curve for each viral target was generated with RNA extracts resulting from the serial dilution of viral stock suspension in distilled water. The slopes (S) of the regression lines were used to calculate the amplification efficiency (E) of the quantitative real-time RT-PCR reactions according to the formula $E=10^{1/s}-1$ to determine the performance of quantitative real-time RT-PCR assays. HAV, HEV, norovirus GI, norovirus GII, and MNV-1 recovery rate percentages from spiked samples were calculated by using the standard curves obtained with viral inoculum dilution and the following formula: quantity of virus recovered after spiking experiments/quantity of viral inoculum X100. The HAV, HEV, norovirus GI or norovirus GII RNA transcript was used as an external amplification control (EAC) to monitor RT-PCR inhibition in samples. This approach has been described in ISO 15216-1, where an external control RNA (i.e. an RNA species carrying the target sequence of interest) is added to an aliquot of RNA sample. The degree of RT-PCR inhibition in each tested sample is obtained by comparing these results with the results of EAC RNA in the absence of sample RNA (i.e. in water).

2.7. Statistical analysis

All statistical analyses were performed using the Statgraphics Centurion XVII software (Statgraphics Centurion Version 17.1.04). The influence of the operator factor on virus recovery rates (norovirus, HAV, HEV and MNV) was first assessed by using one-way analysis of variance (ANOVA). The result of the ANOVA is a p value associated with the hypothesis that the mean recovery rates of all groups were the same. The influence of additional factors on extraction yields of pathogenic viruses (norovirus, HAV, HEV and MNV) was studied by using a one-way ANOVA. The effect of virus (norovirus, HAV, HEV and MNV-1) on virus

229 recovery rates was first evaluated. Because the extraction yields were statistically different
230 according to the virus used (ANOVA, $p < 0.05$), a multiple-comparison procedure (Fisher's
231 least-significant-differences (LSD)) was applied to determine which viruses, provided the
232 highest extraction yields. Graphs plotting the mean and its standard error for each group
233 illustrate the multiple comparison procedure. When confidence intervals for means do not
234 overlap, the difference between two groups of a factor is significant. Next, four factors were
235 tested on extraction yields: (1) dilution of RNA extracts (pure vs. 10-fold diluted), (2) the
236 inoculated level of viruses, (3) setting experiments (R1 to R4) and (4) EAC recovery rates
237 according to the type of food matrix.

238 The limit of detection values (LOD) which correspond respectively to 50% (LOD₅₀) and 95%
239 (LOD₉₅) of the probability of detection were calculated with the method for estimating POD
240 (probability of detection) function and the LOD of a qualitative microbiological measurement
241 method as described by Wilrich et al. (2009). The POD-LOD calculation software was used
242 (version 9, dated 2017-09-23) (Wilrich et al., 2009). This program can be freely downloaded
243 from [www.wiwiwiss.fu-](http://www.wiwiwiss.fu-berlin.de/fachbereich/vwl/iso/ehemalige/professoren/wilrich/index.html)
244 [berlin.de/fachbereich/vwl/iso/ehemalige/professoren/wilrich/index.html](http://www.wiwiwiss.fu-berlin.de/fachbereich/vwl/iso/ehemalige/professoren/wilrich/index.html). The limit of
245 quantification (LOQ) was estimated for each virus using the total error approach based on
246 the accuracy profile (Hubert et al, 2007).

247 The accuracy profile is calculated by a two-sided β expectation tolerance interval (β -ETI) for
248 each level from the results of the validation experiments expressed, such as recovery
249 (trueness) and precision (repeatability and intermediate precision). Data are \log_{10}
250 transformed and we defined the acceptability limits (λ) at ± 95 % of bias because the
251 expected recovery is low. To calculate the accuracy profile a specified proportion has been
252 defined. This proportion is fixed to 80% (β) and corresponds to future measurements would

fall within the acceptability limits. The complete theory and calculation of accuracy profiles are described in the references (Hubert et al, 2007a, Hubert et al, 2007b, Hubert et al, 2008) and a fully developed application is presented for microbiological methods by Feinberg et al (2009) and Boubetra et al (2011). The procedure reported here is simplified and tailored to molecular methods, such as published by Saint-Cyr et al (2014).

The results are presented through a graphical representation as follows: the horizontal (x) axis shows the reference level in log₁₀ concentration level (genome copies/g) and the vertical (y) axis shows at each level the bias, the acceptability limits (λ), and the β -ETI limits.

3. Results

3.1 Mean virus recoveries of norovirus, HAV and HEV according to operator

To determine whether the operator factor influenced virus recovery rates, the mean recovery rates obtained for all settings were compared for each operator (A and B) (Figure 1). These results showed that the mean recovery rates for all settings were 8.04% and 13.81% for HAV, 41.25% and 49.59% for HEV, 9.95% and 6.55% for norovirus GI, 15.78% and 14.88% for norovirus GII, **18.18% and 13.38% for MNV-1** for operator A and operator B respectively. The statistical analysis revealed that the operator factor did not influence virus recovery from multicomponent food (one-way ANOVA; p-value=0.3215).

3.2 Virus recoveries and limit of detection

Table 3 gives the mean extraction yields obtained for norovirus GI, norovirus GII, HAV, HEV and the process control virus according to the inoculum level and repeat experiments (R1 to R4). As expected, no viral RNA was detected in the uninoculated samples. The average of

276 recoveries with pure RNA extracts ranged from 0.35% to 40.44% for HAV, from 5.19% to
277 100% for HEV, from 0.1% to 40.61% for norovirus GI and 0.88% to 69.16% for norovirus GII.

278 The mean recovery rates obtained for HAV, HEV, norovirus GI and norovirus GII showed that
279 the recovery rates from composite matrices vary according to the virus inoculated (one-way
280 ANOVA; p -value<0.001) (Figure 2). More specifically, the multiple comparison tests showed
281 that the recovery rates of HEV were higher and significantly different from other viruses.

282 By testing the 10-fold diluted RNA extracts, the extraction yields obtained for HAV, HEV and
283 norovirus in composite matrices were improved by a factor that ranged from 0.0 to 25.7
284 (Table 3).

285 The process control (MNV-1) was detected in 120 out of 128 RNA extracts analysed and was
286 recovered with an efficiency of between 3.83% and 50.22% for all four food settings.

287 The limits of detection (LOD) for the method were calculated by the Wilrich approach for
288 each virus in each setting with the twelve samples. The LOD₅₀ and LOD₉₅ values for HAV,
289 HEV, norovirus GI and norovirus GII are shown in Table 4.

290 The LOD₉₅ values were 10² genome copies/g for HAV, HEV and norovirus GII and 10³ genome
291 copies/g for norovirus GI. The LOD₅₀ values were between 80 and 92 genome copies/g for
292 HAV, HEV and norovirus GII and 360 genome copies/g for norovirus GI.

293 The limit of quantification (LOQ) was assessed for different viruses using the accuracy
294 profile. Figure 3 shows a typical accuracy profile for HAV. Table 5 summarizes the results for
295 the LOQ and LOD and some performance characteristics (repeatability and intermediate
296 precision). The LOQ was 2.90x10⁴, 1.40x10³, 1.60x10⁴, 1.30x10⁴ genome copies/g for HAV,
297 HEV, norovirus GI and norovirus GII respectively. The LOD for HAV, HEV, norovirus GI and
298 norovirus GII was established at 0.97x10⁴, 0.47x10⁴, 0.53x10⁴ and 0.43x10⁴ genome copies/g
299 respectively.

3.3 Influence of experimental factors on virus extraction yield

To assess the influence of the different experimental factors on HAV, HEV and norovirus extraction from multicomponent matrices, the mean recovery rates from virus-spiked samples were compared. Figure 4 shows the recovery rates for HAV, HEV and norovirus according to their inoculation levels. The statistical analysis indicated that the recovery rates were not significantly different for norovirus GII regardless of the inoculation level (one-way ANOVA; p-value=0.2942 for norovirus GII), but were significantly different for norovirus GI, HAV and HEV when different inoculation levels were tested (one-way ANOVA; p-value<0.001 for HAV and norovirus GI, and p-value<0.0243 for HEV).

More specifically, the multiple-comparison tests showed that inoculations with 2.90×10^3 HAV genome copies, 1.40×10^2 HEV genome copies and 1.6×10^3 norovirus GI genome copies differed significantly from other inoculation levels of HAV, HEV and norovirus GI.

Figure 5 shows the mean recovery rates of HAV, HEV, norovirus in the four repeat experiments. The differences between R1 to R4 were not significant for the extraction yield of norovirus GI (one-way ANOVA; p-value=0.0736) but were significant for the extraction yield of norovirus GII, HAV and HEV (one-way ANOVA; p-value=0.0028 for HAV, one-way ANOVA; p-value<0.001 for norovirus GII and for HEV). More specifically, the multiple-comparison tests showed that R3 was significantly different from R1, R2 and R4 for HAV. Both R1 and R3 were significantly different from R4 for HEV, and R4 was significantly different from R1, R2 and R3 for norovirus GII.

The statistical analysis also revealed that the allocated food setting (factor R1 to R4) influences mean MNV-1 recoveries (one-way ANOVA; p-value=0.0046). The multiple-comparison tests showed that R1 and R2 were significantly different from R3 and R4. Mean

MNV-1 recoveries from allocated food settings R1 and R2 were higher than allocated food settings R3 and R4.

3.4 Recovery rates for external amplification control (EAC)

An EAC corresponding to the viral target was used to examine quantitative real-time RT-PCR inhibition. Table 6 shows the mean percentages of quantitative real-time RT-PCR inhibition for each of the twelve samples. The mean percentages of quantitative real-time RT-PCR inhibition varied from 3.25% to 28.70%. They vary significantly with the type of food matrix (one-way ANOVA; p-value=0.0209) but not with repeated experiments (R1 to R4) (one-way ANOVA; p-value=0.0554).

4. Discussion

Enteric viruses are the leading cause of foodborne outbreaks. Their detection is challenging because many of them are either difficult or impossible to replicate in cell culture, and the number of viral particles present in the food may be very low. Today, quantitative real-time RT-PCR is widely used for virus detection because it is sensitive, specific, rapid and can deliver quantitative data. ISO standards 15216-1 and 15216-2 were published in 2017 and 2019 respectively for detecting and quantifying norovirus and HAV in high-risk food categories such as shellfish, bottled water and vegetables. The method used in this study to recover viruses from composite foods is adapted from the ISO 15216 procedure described for vegetables. The LOD₉₅ varied from 10² to 10³ genome copies per g regardless of the virus analysed. Because of the complexity of the food matrices used in our study, the LOD₉₅ values obtained for norovirus and HAV in composite foods and multicomponent foodstuffs were higher than the LOD₉₅ recently reported for lettuce or raspberries (Lowther et al., 2019).

Derived from the LOQ estimated by accuracy profile, the LOD in our study ranged between 4.33x10³ and 9.67x10³ genome copies per g. These values are slightly different than the estimated LOD₉₅ but the difference, which is around one log, can be explained by the definition of these parameters. LOD₉₅ is the level at which the probability of detection is equal to 95 % (ISO standard 16140-1), whereas the LOD is estimated using a quantitative approach based on a chemical method. Classically, the LOD is the lowest level that differs from the background noise with an accepted confidence level. Armbruster et al. (2008) have previously explained that the LOD is a value as mean + k SD with k, the coverage factor, ranged between 2 and 10. The common value for k is 3 for LOD and if k = 10 it is the LOQ. However, the LOQ is the lowest level quantifiable with a defined trueness and precision (ISO standard 16140-1). In our case, we defined the LOQ using an accuracy profile with a defined trueness and precision, then derived the LOD from this LOQ. These two approaches are totally different because one is for a qualitative approach and the other for a quantitative approach. The results are nonetheless of the same magnitude and consequently the LOD confirms the LOD₉₅ values.

Various methods have been described for extracting viruses from vegetables, composite foods or dairy products (Baert et al., 2008; Blaise-Boisseau et al., 2010; Cheong et al., 2009; Coudray et al., 2013; Dubois et al., 2002; El Sanousy et al., 2013; Hennechart et al., 2017; Hida et al., 2013; Hyeon et al., 2011; Kim et al., 2008; Morillo et al., 2012; Pan et al., 2012; Sanchez et al., 2012; Scherer et al., 2010; Schwab et al., 2000; Stals et al., 2011a, 2011b).

The PEG concentration method was used in our study to concentrate viruses from multicomponent foods. Our results are in agreement with data reported in other studies on food. Virus recovery rates ranged from 0.35% to 40.44% for HAV, 5.19% to 100% for HEV and

0.10% to 69.16% for norovirus. The PEG concentration method was previously developed and applied to recover viruses from suspected food such as shellfish, smooth-surfaced, semi-dried tomatoes or food samples such as spaghetti, chicken, ham and sausages (Martin-Latil et al., 2014, 2012a; Saito et al., 2015; Summa et al., 2012; Rutjes et al., 2006). The norovirus recovery rates in the literature varied between 0.02% and 47%, while the HEV recovery rate from pig liver sausage ranged from 3.94 to 18.38% and the HAV recovery rate from semi-dried tomatoes ranged from 2.91% to 50.92% (Hennechart et al., 2017; Martin-Latil et al., 2014, 2012a; Saito et al., 2015; Summa et al., 2012; Rutjes et al., 2006). Taking into account the mean recovery rates obtained with other matrices, our results lie within the same range.

ISO standard 15216-1 was validated for the detection and quantification of HAV and norovirus in seven food matrices: bottled water, food surfaces, Pacific oysters (*Crassostrea gigas*), common mussels (*Mytilus edulis*), raspberries, lettuce and green onions (Lowther et al., 2017). Other methods for the detection of viruses in semi-dried tomatoes, raw pig livers and dairy products have also been characterised (Hennechart et al., 2017; Martin-Latil et al., 2014, 2012a). However, to our knowledge, no publications until now have reported the validation of a method for detecting norovirus, HAV or HEV from multicomponent foodstuffs.

The experimental design from ISO standard 16140-4:2017 was applied to characterise a method used to detect norovirus, HAV and HEV in multicomponent foodstuffs. In addition to the factors studied in the ISO 16140 standard, part 4 describes the calculation of repeatability and reproducibility. The experimental design allowed us to study a method without a reference method. Twelve different foodstuffs (vegetables, mixed vegetables,

395 meals including meat or fish, soup or sauce) were selected and analysed in this study. This
396 selection allowed us to take into account a wide selection of multicomponent foods
397 representative of **samples analysed** in the laboratory when investigating viral foodborne
398 outbreaks. The items selected are consumed in a variety of forms and are major components
399 of meals.

400 To calculate repeatability, foods were randomly allocated to four different settings (R1, R2,
401 R3 and R4) and results show that the differences between repeated experiments (R1 to R4)
402 were significant for the extraction rate of norovirus GII, HAV and HEV. Different studies have
403 shown that the composition of food products can affect virus extraction (Blaise-Boisseau et
404 al., 2010; Butot et al., 2007; Summa et al., 2012; Yavarmanesh et al., 2013, 2010). **Due to the**
405 **presence of substances that can inhibit PCR amplification, the implementation of different**
406 **controls such as the virus process control and EAC are necessary to validate results.**
407 **According to the recommendations in ISO 15216, the inhibition rates for RNA extracted from**
408 **food samples have to be lower than 75% and virus process control extraction yields higher**
409 **than 1%.** In this study, the mean percentages of **quantitative real-time RT-PCR** inhibition
410 were always lower than 75% and the recovery rate of MNV-1 was more than 1% in 93% of
411 the RNA extracts analysed. In our study, MNV-1 was successfully tested as a process control
412 virus for detecting HAV, HEV and norovirus in multicomponent foodstuffs. Its use had
413 previously been described in the literature for different types of water, shellfish, milk
414 products, meat such as sausages, soft red fruits, lettuces and ready-to-eat foods (Coudray et
415 al., 2013; Hennechart et al., 2015; Martin-Latil et al., 2014; Sanchez et al., 2012; Stals et al.,
416 2011a, 2011b).

417 According to the ISO 16140 procedure, **analyses were conducted** by two operators. An
418 evaluation of operator influence showed no significant difference between the two

operators. The calibration and maintenance of equipment should also limit the difference between operators working at the same laboratory. To evaluate the quality of virus detection from multicomponent foodstuffs for other laboratories, it should be necessary to carry out an inter-laboratory assay.

To conclude, the method described herein could be used to detect viruses in composite food products for routine diagnosis needs. This research underscores the importance of further research to develop, standardise and validate methods for detecting viruses from other food matrices or other viruses than those described in standard ISO 15216.

Acknowledgements

We would like to thank P. Pothier (CNR des virus des gastroentérites, Dijon, France) for providing contaminated stool samples.

We are also grateful to N. Pavio (ANSES, Animal Health Laboratory, Unité Mixte de Recherche (UMR) 1161 Virology, Institut National de la Recherche Agronomique (INRA), École Nationale Vétérinaire d'Alfort (ENVA), Maisons-Alfort, France) for providing HEV-contaminated pig faeces.

References

Anonymous, 2016: NF EN ISO 16140-1: 2016: Microbiology of the food chain-Method validation-Part 1: Vocabulary. International Organization for Standardization, Geneva.

Anonymous, 2018 : PR ISO/DIS 16140-4: 2018-01: Microbiology of the food chain-Method validation-Part 4: Protocol for single-laboratory (in-house) method validation. International Organization for Standardization, Geneva.

443 Anonymous, 2017 EN ISO 15216-1:2017. Microbiology of the food chain–Horizontal method for
 444 determination of hepatitis A virus and norovirus using real-time RT-PCR- Part1: Method for
 445 quantification. International Organization for Standardization, Geneva.

446 Anonymous, 2019 EN ISO 15216-2:2019. Microbiology of the food chain-Horizontal method for
 447 determination of hepatitis A virus and norovirus using real-time RT-PCR- Part2: Method for
 448 detection. International Organization for Standardization, Geneva.

449 Armbruster, D.A., Pry, T. 2008. Limit of blank, limit of detection and limit of quantitation. Clin.
 450 Biochem. Rev. 29, Suppl (i) S49-S52.

451 Baert, L., Uyttendaele, M., Debevere, J. 2008. Evaluation of viral extraction methods on a broad
 452 range of ready-to-eat foods with conventional and real-time RT-PCR for Norovirus GII
 453 detection. Int. J. Food Microbiol. 123, 101-108.

454 Barrabeig, I., Rovira, A., Buesa, J., Bartolome, R., Pinto, R., Prellezo, H., Dominguez, A., 2010.
 455 Foodborne norovirus outbreak: The role of an asymptomatic food handler. BMC Infect. Dis.
 456 10, 269.

457 Bernard, H., Faber, M., Wilking, H., Haller, S., Hohle, M., Schielke, A., Ducomble, T., Siffczyk, C.,
 458 Merbecks, S.S., Fricke, G., Hamouda, O., Stark, K., Werber, D., Outbreak Investigation Team,
 459 2014. Large multistate outbreak of norovirus gastroenteritis associated with frozen
 460 strawberries, Germany, 2012. Euro Surveill. 19, 20719–20727.

461 Blaise-Boisseau, S., Hennechart-Collette, C., Guillier, L., Perelle, S., 2010. Duplex real-time qRT-PCR
 462 for the detection of hepatitis A virus in water and raspberries using the MS2 bacteriophage
 463 as a process control. J. Virol. Methods 166, 48-53.

464 Boubetra, A., Le Nestour, F., Allaert, C., Feinberg, M., 2011. Analysis of drinking water and their
 465 validation of alternative methods for their application to escherichia coli. Appl. Environ.
 466 Microbiol. 77, 3360–3367.

467 Butot, S., Putallaz, T., Sanchez, G., 2007. Procedure for rapid concentration and detection of enteric
 468 viruses from berries and vegetables. Appl. Environ. Microbiol. 73, 186-92.

469 Cannon, J.L., Papafragkou, E., Park, G.W., Osborne, J., Jaykus, L.A., Vinje, J., 2006. Surrogates for the
 470 study of norovirus stability and inactivation in the environment: comparison of murine
 471 norovirus and feline calicivirus. *J. Food Prot.* 69, 2761-2765.

472 Cheng, D., Zou, S., Liao, N., Shi, X., Chen, J., Zhang, Y., Sun, L., Zhang, R., 2017. Evaluation of an
 473 extraction method for the detection of GI and GII Noroviruses in fruit and vegetable salads.
 474 *Food Sci.* 83, 393-400.

475 Cheong, S., Lee, C., Choi, W.C., Lee, C.H., Kim, S.J., 2009. Concentration method for the detection of
 476 enteric viruses from large volumes of foods. *J. Food Prot.* 72, 2001–2005.

477 Costafreda, M. I., Bosch, A., Pinto, R. M., 2006. Development, evaluation, and standardization of a
 478 real-time taqman reverse transcription-Pcr assay for quantification of Hepatitis A virus in
 479 clinical and shellfish samples. *Appl. Environ. Microbiol.* 72, 3846-55.

480 Coudray, C., Merle, G., Martin-Latil, S., Guillier, L., Perelle, S., 2013. Comparison of two extraction
 481 methods for the detection of hepatitis A virus in lettuces using the murine norovirus as a
 482 process control. *J. Virol. Methods*, 193, 96-102.

483 Cromeans, T., Sobsey, M.D., Fields, H.A., 1987. Development of a plaque assay for a cytopathic,
 484 rapidly replicating isolate of hepatitis A virus. *J. Med. Virol.* 22, 45-56.

485 Dalton, H.R., Hunter, J.G., Bendall, R.P. 2013. Hepatitis E. *Current opinion in infectious diseases*, 26,
 486 471-478.

487 da Silva, A.K., Le Saux, J.C., Parnaudeau, S., Pommepuy, M., Elimelech, M., Le Guyader, F.S., 2007.
 488 Evaluation of removal of noroviruses during wastewater treatment, using real-time reverse
 489 transcription-PCR: different behaviors of genogroups I and II. *Appl. Environ. Microbiol.* 73,
 490 7891-7897.

491 Donnan, E.J., Fielding, J.E., Gregory J.E., Lalor, K., Rowe, S., Goldsmith, P., Antoniou, M., Fullerton,
 492 K.E., Knope, K., Copland, J.G., Bowden, D.S., Tracy, S.L., Hogg, G.G., Tan, A., Adamopoulos, J.,
 493 Gaston, J., Vally, H., 2012. A multistate outbreak of hepatitis A associated with semidried
 494 tomatoes in Australia, 2009. *Clin. Infect. Dis.* 54.

495 Dubois, E., Agier, C., Traore, O., Hennechart, C., Merle, G., Cruciere, C., Laveran, H. 2002. Modified
 496 concentration method for the detection of enteric viruses on fruits and vegetables by reverse
 497 transcriptase-polymerase chain reaction or cell culture. *J. Food Prot.* 65, 12, 1962-1969.

498 El-Senousy, W.M., Costafreda, M.I., Pintó, R.M., Bosch, A. 2013. Method validation for norovirus
 499 detection in naturally contaminated irrigation water and fresh produce. *Int.J. Food Microbiol.*
 500 167, 74-79.

501 Ethelberg, S., Lisby, M., Bottiger, B., Schultz, A.C., Villif, A., Jensen, T., Olsen, K.E., Scheutz, F., Kjelso,
 502 C., Muller, L., 2010. Outbreaks of gastroenteritis linked to lettuce, Denmark, January 2010.
 503 *Euro Surveill.* Feb 11, 15, 6.

504 Feinberg, M., Sohler, D., David, J.-F., 2009. Validation of an alternative method for counting
 505 enterobacteriaceae in foods based on accuracy profile. *J. AOAC Int.* 92, 527–537.

506 Fournet, N., Baas, D., van Pelt, W., Swaan, C., Ober, H., Isken, L., Cremer, J., Friesema, I., Vennema,
 507 H., Boxman, I., Koopmans, M., Verhoef, L., 2012. Another possible foodborne outbreak of
 508 hepatitis A in the Netherlands indicated by two closely related molecular sequences, July to
 509 October 2011. *Euro Surveill.* 17, 20079–20081.

510 Fraisse, A., Coudray-Meunier, C., Martin-Latil, S., Hennechart-Collette, C., Delannoy, S., Fach, P.,
 511 Perelle, S., 2017. Digital RT-PCR method for hepatitis A virus and norovirus quantification in
 512 soft berries. *Int. J. Food Microbiol.* 21, 243, 36-45.

513 Franck, K.T., Lisby, M., Fonager, J., Schultz, A.C., Bottiger, B., Villif, A., Absalonsen, H., Ethelberg, S.
 514 2015. Sources of Calicivirus contamination in foodborne outbreaks in Denmark, 2005-2011-
 515 the role of the asymptomatic food handler. *J. I. Dis.* 211, 563-570.

516 Gallot, C., Grout, L., Roque-Afonso, A.M., Couturier, E., Carrillo-Santistevé, P., Pouey, J., Letort, M.J.,
 517 Hoppe, S., Capdepon, P., Saint-Martin, S., De Valk, H., Vaillant, V., 2011. Hepatitis A
 518 associated with semidried tomatoes, France, 2010. *Emerg. Infect. Dis.* 17, 566-567.

519 Hamza, I.A., Jurzik, L., Uberla, K., Wilhelm, M. (2011). Methods to detect infectious human enteric
 520 viruses in environmental water samples. *Int. J. Hyg. Environ. Health* 214, 424-436.

521 Hennechart-Collette, C., Niveau F., Martin-Latil S., Fraisse A., Perelle S., 2019. Development of an
 522 extraction method to detect enteric viruses in dressed vegetables. *Int. J. Food Microbiol.*
 523 311:108349.

524 Hennechart-Collette, C., Martin-Latil, S., Fraisse, A., Perelle, S., 2017. Comparison of three extraction
 525 methods to detect noroviruses in dairy products. *Food Microbiol.* 61, 113-119.

526 Hennechart-Collette, C., Martin-Latil, S., Guillier, L., Perelle, S., 2015. Determination of which virus to
 527 use as a process control when testing for the presence of hepatitis A virus and norovirus in
 528 food and water. *Int. J. Food Microbiol.* 202, 57-65.

529 Hennechart-Collette, C., Martin-Latil, S., Guillier, L., Perelle, S., 2014. Multiplex real-time RT-qPCR for
 530 the detection of Norovirus in bottled and tap water using murine norovirus as a process
 531 control. *J. Appl. Microbiol.* 116, 179-190.

532 Herman, K.M., Hall, A.J., Gould, L.H., 2015. Outbreaks attributed to fresh leafy vegetables, United
 533 States, 1973-2012. *Epidemiol. Infect.* 143, 14, 3011-3021.

534 Hida, K., Kulka, M., Papafragkou, E., 2013. Development of a rapid total nucleic acid extraction
 535 method for the isolation of hepatitis A virus from fresh produce. *Int. J. Food Microbiol.*, 161,
 536 143-150.

537 Hubert, P., Nguyen-Huu, J.J., Boulanger, B., Chapuzet, E., Chiap, P., Cohen, N., Compagnon, P.A.,
 538 Dewé, W., Feinberg, M., Lallier, M., Laurentie, M., Mercier, N., Muzard, G., Nivet, C., Valat,
 539 L., & Rozet, E., 2007a. Harmonization of strategies for the validation of quantitative analytical
 540 procedures a SFSTP proposal – Part II. *J. Pharm. Biomed. Anal.* 45, 70–81.

541 Hubert, P., Nguyen-Huu, J.J., Boulanger, B., Chapuzet, E., Cohen, N., Compagnon, P.A., Dewé, W.,
 542 Feinberg, M., Laurentie, M., Mercier, N., Muzard, G., Valat, L., & Rozet, E., 2007b,
 543 Harmonization of strategies for the validation of quantitative analytical procedures a SFSTP proposal –
 544 Part III 2007. *J. Pharm. Biomed. Anal.* 45, 82–96.

545 Hubert, P., Nguyen-Huu, J.J., Boulanger, B., Chapuzet, E., Cohen, N., Compagnon, P.A., Dewé, W.,
 546 Feinberg, M., Laurentie, M., Mercier, N., Muzard, G., Valat, L., & Rozet, E., 2008.

Harmonization of strategies for the validation of quantitative analytical procedures: A SFSTP proposal. Part IV. Examples of application . J. Pharm. Biomed. Anal. 48, 760–771.

Hyeon, J.Y., Chon, J.W., Park, C., Lee, J.B., Choi, I.S., Kim, M.S., Seo, K.H., 2011. Rapid detection method for hepatitis A virus from lettuce by a combination of filtration and integrated cell culture-real-time reverse transcription PCR. J. Food Prot. 74, 1756-1761.

Kim, H.Y., Kwak, I.S., Hwang, I.G., Ko, G., 2008. Optimization of methods for detecting norovirus on various fruit. Journal of virological methods, 153, 104-110.

Jothikumar, N., Lowther, J.A., Henshilwood, K., Lees, D.N., Hill, V.R., Vinje, J., 2005. Rapid and sensitive detection of noroviruses by using TaqMan-based one-step reverse transcription-PCR assays and application to naturally contaminated shellfish samples. Appl. Environ. Microbiol. 71, 1870-1875.

Kageyama, T., Kojima, S., Shinohara, M., Uchida, K., Fukushi, S., Hoshino, F.B., Takeda, N., Katayama, K., 2003. Broadly reactive and highly sensitive assay for Norwalk-like viruses based on real-time quantitative reverse transcription-PCR. J. Clin. Microbiol. 41, 1548-1557.

Kotwal, G. and Cannon, J.L., 2014. Environmental persistence and transfer of enteric viruses. Curr. Opin .Virol. 4, 37-43.

Le Guyader, F.S., Krol, J., Ambert-Balay, K., Ruvoen-Clouet, N., Desaubliaux, B., Parnaudeau, S., Le Saux, J.C., Ponge, A., Pothier, P., Atmar, R.L., Le Pendu, J., 2010. Comprehensive analysis of a norovirus-associated gastroenteritis outbreak, from the environment to the consumer. J. Clin. Microbiol. 48, 915–920.

Loisy, F., Atmar, R.L., Guillon, P., Le Cann, P., Pommepuy, M., Le Guyader, F.S., 2005. Real-time RT-PCR for norovirus screening in shellfish. J. Methods. 123, 1-7.

Lowther, J.A, Bosch, A., Butot, S., Ollivier, J., Mäde, D., Rutjes, S.A., Hardouin, G., Lombard, B., in't Veld, P., Leclercq, A., 2017. Validation of ISO 15216 part 1 –Quantification of hepatitis A virus and norovirus in food matrices. Int. J. Food Microbiol. 288, 82-90.

572 Martin-Latil, S., Hennechart-Collette, C., Guillier, L., Perelle, S., 2014. Method for HEV detection in
 573 raw pig liver products and its implementation for naturally contaminated food. *International*
 574 *J. Food Microbiol.* 176, 1-8.

575 Martin-Latil, S., Hennechart-Collette, C., Guillier, L., Perelle, S., 2012a. Comparison of two extraction
 576 methods for the detection of hepatitis A virus in semi-dried tomatoes and murine norovirus
 577 as a process control by duplex RT-qPCR. *Food Microbiol.* 31, 246-253.

578 Martin-Latil, S., Hennechart-Collette, C., Guillier, L., Perelle, S., 2012b. Duplex RT-qPCR for the
 579 detection of hepatitis E virus in water, using a process control. *International J. Food*
 580 *Microbiol.* 157, 167-173.

581 Matthews, J.E., Dickey, B.W., Miller, R.D., Felzer, J.R., Dawson, B.P., Lee, A.S., Rocks, J.J., Kiel, J.,
 582 Montes, J.S., Moe, C.L., et al., 2012. The epidemiology of published norovirus outbreaks: a
 583 review of risk factors associated with attack rate and genogroup. *Epidemiol Infect.* 140, 1161-
 584 1172.

585 Meng, X.J., 2010. Hepatitis E virus: animal reservoirs and zoonotic risk. *Vete. Microbiol.* 140, 256-265.

586 Morillo, S.G., Luchs, A., Cilli, A., do Carmo Sampaio Tavares Timenetsky, M., 2012. Rapid detection of
 587 norovirus in naturally contaminated food: foodborne gastroenteritis outbreak on a cruise
 588 ship in Brazil, 2010. *Food Environ. Virol.* 4, 3, 124-129.

589 Müller, L., Rasmussen, L.D., Jensen, T., Schultz, A.C., Kjelsø, C., Barnadas, C., Sigsgaard, K., Larsen,
 590 A.R., Widstrup Jensen, C., Jeppesen, S., Uhrbrand, K., Hove, N., Mølbak, K., Ethelberg, S.,
 591 2016. Series of norovirus outbreaks caused by consumption of green coral lettuce, Denmark,
 592 April 2016. *PLoS Curr.* 4, 8.

593 Lemon, S.M., Murphy, P.C., Shields, P.A., Ping, L.H., Feinstone, S.M., Cromeans, T., Jansen, R.W.,
 594 1991. Antigenic and genetic variation in cytopathic hepatitis A virus variants arising during
 595 persistent infection: evidence for genetic recombination. *J. Virol.* 65, 2056e2065.

596 Pan, L., Zhang, Q., Li, X., Tian, P., 2012. Detection of human norovirus in cherry tomatoes, blueberries
 597 and vegetable salad by using a receptor-binding capture and magnetic sequestration
 598 (RBCMS) method. *Food Microbiol.* 30, 420-426.

599 Pavio, N., Lunazzi, A., Barnaud, E., Bouquet, J., Rogée, S., 2010. Hépatite E: nouvelles connaissances.
 600 *Bull.Epid. N°38/Sécial zoonoses.*

601 Pavio, N., Renou, C., Boutrouille, A., Eloit, M., 2006. L'hépatite E : une zoonose méconnue. *Virol.* 10,
 602 341-351.

603 Pinto, R.M., Costafreda, M.I., Bosch, A., 2009. Risk assessment in shellfish-borne outbreaks of
 604 hepatitis A. *Appl. Environ. Microbiol.* 75, 7350-7355.

605 Rönqvist, M., Aho, E., Mikkela, A., Ranta, J., Tuominen, P., Ratto, M., Maunula, L., 2014. Norovirus
 606 transmission between hands, gloves, utensils, and fresh produce during simulated food
 607 handling. *Appl.Environ. Microbiol.* 80, 5403-5410.

608 Rutjes, S.A., van den Berg, H.H., Lodder, W.J., de Roda Husman, A.M., 2006. Real-time detection of
 609 noroviruses in surface water by use of a broadly reactive nucleic acid sequence-based
 610 amplification assay. *Appl. Environ. Microbiol.* 72, 8, 5349-5358.

611 Saito, H, Toho, M, Tanaka, T, Noda, M., 2015. Development of a practical method to detect
 612 Noroviruses contamination in composite meals food. *Environ. Virol.* 7, 3, 239-248.

613 Saint-Cyr, M.J., Perrin-Guyomard, A., Houée, P., Vasseur, M.V., Laurentie M., 2014. Use of accuracy
 614 profile procedure to validate a real-time PCR method to quantify bacteria in feces. *JAOAC*
 615 *Int.* 97, 2, 1-7.

616 Sanchez, G., Elizaquivel, P., Aznar, R., 2012. A single method for recovery and concentration of
 617 enteric viruses and bacteria from fresh-cut vegetables. *Int. J. Food Microbiol.* 152, 9-13.

618 Santé publique France, 2017. Données relatives aux toxi-infections alimentaires collectives déclarées
 619 en France en 2017 (Data on collective food-borne outbreaks reported in France in 2017).
 620 Sante Publique France. Available from: <https://www.santepubliquefrance.fr/maladies-et->

621 traumatismes/maladies-infectieuses-d-origine-alimentaire/toxi-infections-alimentaires-
622 collectives/documents/bulletin-national/donnees-relatives-aux-toxi-infections-alimentaires-
623 collectives-declarees-en-france-en-2017.

624 Sarvikivi, E., Roivainen, M., Maunula, L., Niskanen, T., Korhonen, T., Lappalainen, M., Kuusi, M., 2012.
625 Multiple norovirus outbreaks linked to imported frozen raspberries. *Epidemiol. Infect.* 140,
626 260–267.

627 Scherer, K., Johne, R., Schrader, C., Ellerbroek, L., Jörg Schulenburg, J., Klein, G., 2010. Comparison of
628 two extraction methods for viruses in food and application in a norovirus gastroenteritis
629 outbreak. *J. Virol. Methods.* 169 , 22-27.

630 Schwab, K.J., Neill, F.H., Fankhauser, R.L., Daniels, N.A., Monroe, S.S., Bergmire-Sweat, D.A., Estes,
631 M.K., Atmar, R.L., 2000. Development of methods to detect "Norwalk-like viruses" (NLVs) and
632 hepatitis A virus in delicatessen foods: application to a food-borne NLV outbreak. *Appl.*
633 *Environ. Microbiol.* 66, 1, 213-218.

634 Stals, A., Uyttendaele, M., Baert, L., Van Coillie, E., 2013. Norovirus transfer between foods and food
635 contact materials. *Journal of food protection*, 76, 1202-1209.

636 Stals, A., Baert, L., De Keuckelaere, A., Van Coillie, E., Uyttendaele, M., 2011a. Evaluation of a
637 norovirus detection methodology for ready-to-eat foods. *Int. J.Food Microbiol.* 145, 420-425.

638 Stals, A., Baert, L., Van Coillie, E., Uyttendaele, M., 2011b. Evaluation of a norovirus detection
639 methodology for soft red fruits. *Food Microbiol.* 28, 52-58.

640 Summa, M., von Bonsdorff, C.H., Maunula, L., 2012. Evaluation of four virus recovery methods for
641 detecting noroviruses on fresh lettuce, sliced ham, and frozen raspberries. *J. Virol.Methods.*
642 183, 154-160.

643 Svraka, S., Duizer, E., Vennema, H., de Bruin, E., van der Veer, B., Dorresteyn, B., Koopmans, M.,
644 2007. Etiological role of viruses in outbreaks of acute gastroenteritis in the Netherlands from
645 1994 through 2005. *J. Clin.Microbiol.* 45, 1389-1394.

646 Thebault, A., Teunis, P.F., Le Pendu, J., Le Guyader, F.S., Denis, J.B., 2013. Infectivity of GI and GII
 647 noroviruses established from oyster related outbreaks. *Epidemics* 5, 98–110.

648 Yavarmanesh, M., Abbaszadegan, M., Alum, A., Mortazavi, A., Habibi Najafi, M. B., Bassami, M. R.,
 649 Nassiri, M. R., 2013. Impact of milk components on recovery of viral RNA From MS2
 650 bacteriophage. *Food and Env. Virol.* 5, 103-9.

651 Yavarmanesh, M., Abbaszadegan, M., Mortazavi, A., Najafi, M. B., Bassami, M. R., Nassiri, M. R.,
 652 2010. Impact of milk components in recovery of the MS2 bacteriophage as an indicator of
 653 enteric viruses. *J. Virol. Methods.* 168, 103-107.

654 Wadl, M., Scherer, K., Nielsen, S., Diedrich, S., Ellerbroek, L., Frank, C., Gatzler, R., Hoehne, M., Johne,
 655 R., Klein, G., et al., 2010. Food-borne norovirus-outbreak at a military base, Germany, 2009.
 656 *BMC Infect. Dis.* 10, 30.

657 Wilrich, C., Wilrich, P.T., 2009. Estimation of the POD function and the LOD of a qualitative
 658 microbiological measurement method. *J. AOAC Int.* 92, 1763-1772.

659
 660
 661




Figure 1: Comparison of mean recovery rates for HAV, HEV, norovirus GI, norovirus GII and MNV according to the operator factor (operator A and operator B). For each virus, 16 analyses were performed by each operator (four foodstuffs for all settings with all levels of inoculation) and RNA extracts were analysed in duplicate (pure and ten-fold diluted) with a quantitative real-time RT-PCR assay




Figure 2: Comparison of mean virus recovery rates from spiked samples for all settings




Figure 3: Example of accuracy profile for HAV. The black line is the bias and blue lines are the tolerance limits that define the 80% tolerance interval around the bias. Yellow points are the concentration levels tested. Dotted lines are the acceptance limit.









Figure 4: Recovery rates for HAV, HEV and norovirus with respect to artificial inoculation levels


HAV



norovirus GI




HEV



Repeat experiments

norovirus GII



Repeat experiments

Figure 5: Recovery rates for HAV, HEV and norovirus in repeat experiments (R1 to R4)

Table 1: Selected food samples according to the matrix type and processing factor

Processing factor	Matrix type	Matrix (EFSA FoodEx2 classification)	Sample number
Tin can	Composite dishes with meat	Meat stew (A03VY)	1
		Vol-au-vent (A040G)	2
	Vegetables	Spinach (A00MH)	3
		Corn salad (A05EA)	4
		French beans (A00PG)	5
	Mixed vegetables	Ratatouille (A03YH)	6
	Soup	Vegetable soup (A041S)	7
	Sauce	Ketchup (A044P)	8
Chopped	Composite dishes with meat or fish	Spaghetti bolognese (A040V)	9
		Paella (A041D)	10
		Mixed salmon and spinach (A0C75 and A00MH)	11
Frozen	Mixed vegetables	Mushrooms (A03YV)	12

Table 2: Experimental design for detection of each enteric virus (HAV, HEV and norovirus) in multicomponent foodstuffs adapted from the ISO standard 16140-4:2017 procedure performed by two operators

Virus contamination levels	Repeat experiments				
	R1	R2	R3	R4	
	Very low	Sample 8	Sample 4	Sample 5	Sample 6
	Low	Sample 1	Sample 11	Sample 6	Sample 3
	Medium	Sample 2	Sample 10	Sample 7	Sample 5
	High	Sample 12	Sample 9	Sample 8	Sample 4

Table 3: Mean percentage recovery calculated for four inoculum levels of HAV, HEV, norovirus GI or GII in the presence of MNV-1

Virus	Number of genome copies	RNA extracts	Repeated experiment R1 (%±SD)	(F)	Repeated experiment R2 (%±SD)	(F)	Repeated experiment R3 (%±SD)	(F)	Repeated experiment R4 (%±SD)	(F)
HAV	2.90x10 ⁶	pure	1.56±1.31 (4/4)	1.6	2.80±1.05 (4/4)	1.2	32.53±3.99 (4/4)	1.5	4.39±0.72 (4/4)	0.8
		10-fold diluted	2.58±2.54 (4/4)		3.28±1.80 (4/4)		48.81±3.96 (4/4)		3.50±0.80 (4/4)	
	2.90x10 ⁵	Pure	3.72±4.01 (4/4)	1.5	3.62±2.34 (4/4)	1.1	7.26±8.27 (4/4)	1.7	0.41±0.67 (4/4)	
		10-fold diluted	5.57±4.52 (4/4)		4.05±2.91 (4/4)		12.16±2.18 (3/4)		nd	
	2.90x10⁴	Pure	3.11±3.49 (4/4)	1.5	3.45±3.36 (4/4)	0.9	4.75±4.06 (4/4)	0.9	4.02±2.54 (4/4)	1.0
		10-fold diluted	4.56±5.29 (3/4)		3.21±2.67 (4/4)		4.33±6.70 (2/4)		0.40±0.63 (4/4)	
	2.90x10 ³	Pure	24.99±21.17 (4/4)	2.3	19.06±16.29 (3/4)	1.2	40.44±14.48 (2/4)	0.1	0.35 (1/4)	
		10-fold diluted	57.90±49.40 (4/4)		22.55 (1/4)		3.13±0.43 (2/4)		nd	
	MNV		8.49±9.84		8.12±6.22		12.28±17.52		3.83±4.85	
	Total samples with recovery rates >1%				27/32					
Virus	Number of genome copies	RNA extracts	Repeated experiment R1 (%±SD)	(F)	Repeated experiment R2 (%±SD)	(F)	Repeated experiment R3 (%±SD)	(F)	Repeated experiment R4 (%±SD)	(F)
HEV	1.40x10⁵	pure	29.14±13.87 (4/4)	2.1	6.24±4.92 (4/4)	2.7	59.04±9.99 (4/4)	1.7	16.65±18.31 (4/4)	0.3
		10-fold diluted	60.22±40.64 (4/4)		16.92±10.99 (4/4)		100.00±0.00 (4/4)		4.62±2.21 (3/4)	
	1.40x10 ⁴	pure	33.64±45.25 (4/4)	2.8	57.34±38.49 (3/4)	1.0	27.49±16.87 (4/4)	2.2	5.19±6.09 (2/4)	0.9
		10-fold diluted	94.45 (1/4)		57.63±51.29 (2/4)		59.87±14.20 (2/4)		4.87±5.96 (3/4)	
	1.40x10 ³	pure	16.09±11.75 (2/4)	6.2	17.90 (1/4)	11.1	100.00±0.00 (3/4)		nd	
		10-fold diluted	100.00±0.00 (3/4)		99.67±99.50 (2/4)		nd		14.62±10.48 (3/4)	
	1.40x10 ²	pure	55.79±62.52 (2/4)		11.64 (1/4)	8.6	100.00±0.00 (3/4)		50.64±53.76 (2/4)	1.7
		10-fold diluted	nd		100.00±0.00 (2/4)		nd		86.35	
MNV-1	MNV		18.43±11.74		15.40±16.00		12.78±8.44		12.33±5.39	
	Total samples with recovery rates >1%				30/32					

Virus	Number of genome copies	RNA extracts	Repeated experiment R1 (%±SD)	(F)	Repeated experiment R2 (%±SD)	(F)	Repeated experiment R3 (%±SD)	(F)	Repeated experiment R4 (%±SD)	(F)
norovirus GI	1.60x10 ⁶	pure	0.10±0.08 (4/4)	0.6	1.77±0.12 (4/4)	0.5	10.95±6.63 (4/4)	1.5	1.42±0.41 (4/4)	1.3
		10-fold diluted	0.06±0.05 (2/4)		0.93±0.36 (4/4)		16.29±11.10 (4/4)		1.91±0.63 (4/4)	
	1.60x10 ⁵	pure	3.36±0.64 (4/4)	1.4	8.24±2.95 (4/4)	0.4	5.59±2.79 (4/4)	2.2	5.44±0.96 (4/4)	0.9
		10-fold diluted	0.48 (1/4)		2.96±3.33 (3/4)		12.01±10.46 (3/4)		5.18±4.23 (3/4)	
	1.60x10 ⁴	pure	2.55±3.34 (2/4)	4.6	nd		3.89±3.14 (4/4)	25.7	11.68±9.93 (3/4)	0.1
		10-fold diluted	11.96 (1/4)		9.76 (1/4)		100.00 (1/4)		1.68 (1/4)	
	1.60x10 ³	pure	1.14±0.54 (2/4)		nd		40.61 (1/4)		nd	
		10-fold diluted	nd		nd		nd		100.00 (1/4)	
MNV-1	MNV		11.99±12.50		14.75±12.65		9.38±8.34		15.49±8.83	
	Total samples with recovery rates >1%				31/32					

Virus	Number of genome copies	RNA extracts	Repeated experiment R1 (%±SD)	(F)	Repeated experiment R2 (%±SD)	(F)	Repeated experiment R3 (%±SD)	(F)	Repeated experiment R4 (%±SD)	(F)
norovirus GII	1.30x10 ⁶	pure	1.85±0.67 (4/4)	1.0	2.08±1.59 (4/4)	1.0	10.16±6.97 (3/4)	1.8	21.63±1.76 (4/4)	1.5
		10-fold diluted	1.78±0.79 (4/4)		2.09±2.09 (4/4)		18.60±4.04 (4/4)		32.06±2.60 (4/4)	
	1.30x10 ⁵	pure	8.10±1.89 (4/4)	0.4	7.98±1.47 (4/4)	0.4	14.24±0.79 (4/4)	0.9	49.94±12.26 (4/4)	1.2
		10-fold diluted	3.39±2.35 (4/4)		3.46±4.11 (4/4)		12.95±3.93 (3/4)		58.92±17.38 (4/4)	
	1.30x10 ⁴	pure	10.32±1.70 (4/4)	0.0	0.88±0.63 (4/4)		8.04±6.01 (4/4)	0.1	69.16±6.90 (4/4)	0.04
		10-fold diluted	0.34 (1/4)		nd		0.37 (1/4)		2.56±2.06 (3/4)	
	1.30x10 ³	pure	4.35 (1/4)		nd		8.89±8.67 (3/4)		nd	
		10-fold diluted	nd		nd		nd		nd	
MNV-1	MNV		50.22±41.50		35.41±35.44		12.08±4.45		11.84±5.04	
	Total samples with recovery rates >1%				32/32					

The mean of operator A and B replicates was used for each inoculation level sample. Results are expressed as means +/- SD. The number of positive Ct determinations is mentioned for HAV, HEV, norovirus GI and GII. RNA extracts were tested twice for each operator, resulting in four cycle threshold (Ct) values for each sample. The ratio between the mean values for extraction yields obtained with undiluted RNA extracts and those obtained with 10-fold diluted RNA extracts were calculated to determine whether the dilution of RNA extracts enhanced mean extraction yields (F).

nd: not detected.

Table 4: LOD₅₀ and LOD₉₅ calculated by Wilrich approach for HAV, HEV and norovirus

Virus target	Genome copies/g	R1	R2	R3	R4	All settings
HAV	LOD ₅₀	<116*	6.00x10 ¹	1.16 x10 ²	2.08x10 ²	8.00 x10 ¹
	LOD ₉₅	<116*	2.52x10 ²	4.80 x10 ²	9.20x10 ²	3.52x10 ²
HEV	LOD ₅₀	3.00x10 ¹	1.84x10 ²	1.37x10 ¹	1.60x10 ²	9.20x10 ¹
	LOD ₉₅	1.28 x10 ²	8.00x10 ²	6.00x10 ¹	6.80x10 ²	3.96x10 ²
norovirus GI	LOD ₅₀	3.44x10 ²	1.24x10 ³	1.16x10 ²	2.72x10 ²	3.60x10 ²
	LOD ₉₅	1.48x10 ³	5.20x10 ³	4.80x10 ²	1.16x10 ³	1.56x10 ³
norovirus GII	LOD ₅₀	1.16x10 ²	1.52x10 ²	2.60x10 ¹	1.52x10 ²	9.20x10 ¹
	LOD ₉₅	4.00x10 ²	6.40x10 ²	1.16x10 ²	6.40x10 ²	4.00x10 ²

(*LOD₅₀ and LOD₉₅ values were estimated. It was not possible to determine values because the LOD was not reached)

Table 5: performance characteristics for all viruses

	Virus			
	HAV	HEV	norovirus GI	norovirus GII
LOQ (genome copies/g)	2.90x10 ⁴	1.40x10 ³	1.60x10 ⁴	1.30x10 ⁴
LOD (genome copies/g)	0.97x10 ⁴	0.47x10 ⁴	0.53x10 ⁴	0.43x10 ⁴
Validated range level (genome copies/g)	2.90x10 ⁴ to 2.90x10 ⁶	1.40x10 ³ to 1.40x10 ⁵	1.60x10 ⁴ to 1.60x10 ⁶	1.30x10 ⁴ to 1.30x10 ⁶
Repeatability (genome copies/g)	0.08-0.66	0.48-0.58	0.06-0.59	0.03-0.29
Intermediate precision (genome copies/g)	0.61-1.17	0.69-0.75	0.20-0.86	0.45-0.89

Table 6: Mean percentages of **quantitative real-time RT-PCR** inhibition for each sample

Sample number	Mean inhibition recovery rates (% \pm SD)
1	15.71 \pm 8.56 (N= 8)
2	6.44 \pm 9.15 (N= 8)
3	3.25 \pm 5.77 (N= 8)
4	8.75 \pm 9.29 (N= 16)
5	12.99 \pm 23.26 (N= 16)
6	13.67 \pm 25.95 (N= 16)
7	6.27 \pm 11.60 (N= 8)
8	28.70 \pm 27.00 (N= 16)
9	22.00 \pm 13.36 (N= 8)
10	16.51 \pm 10.33 (N= 8)
11	15.44 \pm 13.08 (N= 8)
12	26.84 \pm 13.39 (N= 8)