



HAL
open science

Characterization of the genetic basis of resistance to (bio)pesticides in *Cydia pomonella*

Mathilda Idier, Myriam Siegwart, Benoit Barrès, Claire Lavigne, Bertrand
Gauffre

► **To cite this version:**

Mathilda Idier, Myriam Siegwart, Benoit Barrès, Claire Lavigne, Bertrand Gauffre. Characterization of the genetic basis of resistance to (bio)pesticides in *Cydia pomonella*. Congrès de la Société Européenne de Biologie Evolutive, Aug 2022, Prague, Czech Republic. anses-04542160

HAL Id: anses-04542160

<https://anses.hal.science/anses-04542160>

Submitted on 11 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

CHARACTERIZATION OF THE GENETIC BASES OF RESISTANCE TO (BIO)-PESTICIDES IN *CYDIA POMONELLA* (L.)

Mathilda IDIER^{1,2}, Myriam SIEGWART¹, Benoit BARRÈS², Claire LAVIGNE¹ and Bertrand GAUFFRE¹


¹ UR1115 Plantes et Systèmes de culture Horticoles, INRAE, Avignon, France

² Université de Lyon, Anses INRAE, USC CASPER, Lyon, France

BACKGROUND

Cydia pomonella (Lepidoptera) is the major pest in apple orchards worldwide

Emergence of resistances in field



Current methods of detection have drawbacks: cost, accuracy, duration (~1 year)

Need to develop more effective tools!

Known types of resistance in *C. pomonella*:

Target-site resistance

Simple and known

Detoxification

Complex and mostly unknown


What are the mechanisms involved in multi-resistance?

Are there specific or common mechanisms depending on the active substance?

Is detoxification the main mechanism at work?

MAIN FINDINGS

A) Corrected mortality (%) for each generation of selection of each of the strains created (repetitions pooled).



B) LD50 of active substances for the 9 sub-strains obtained and the reference susceptible strain. Significant effects compared to the susceptible strain values are noted with *, ** and ***.

Active substances	Strains			Susceptible strain
	F9-10 No exposure strain	F9-10 Medium dose strain	F9-10 High dose strain	
Spinosad	5.66 ns (±2.11)	7.53 *** (±1.288)	7.35 ** (±1.671)	2.84 (±0.963)
Deltamethrin	0.064 ns (±0.025)	0.111 ns (±0.036)	0.193 ns (±0.92)	0.083 (±0.027)
Chlorantraniliprole	0.517 ns (±0.053)	0.823 ns (±0.121)	0.75 ns (±0.111)	0.687 (±0.127)

CONCLUSIONS


- Precise phenotype characterization is complex. → Inconsistency between an effective selection in rearing and phenotype measurements.
- Rather than cross-resistance, it seems that an accumulation of mechanisms conferred resistance to our population, leading to **multiple resistance**.
- Detoxification is not the main mechanism at work, **other mechanisms of metabolic resistance** and **genes involved in other functions** are involved in resistance.
- Several markers were identified as linked to resistance phenotypes, providing a basis for the development of **new biomolecular tools for resistance monitoring** in *C. pomonella*.

METHODS

One population sampled in the South-East of France, characterized as multi-resistant to different insecticides.

It consists of several steps:

A) Selecting pesticide resistant strains through an experimental evolution



3 strains were selected: Spinosad-resistant (SPI), Deltamethrin-resistant (DEL) and Chlorantraniliprole-resistant (CHL). Within each strain, a gradient of dose was applied: no exposure (C), medium dose (M-LD) and high dose (H-LD).


B) Phenotyping using bioassays to assess the resistance level of the strains

C) RNA-pool seq and DNA-pool seq of the F9 to identify the differential expression between reference genome and strains and variants through a filtering approach



$$F_{SNP(\text{control})} \leq F_{SNP(\text{F9-10 no exposure})} \leq F_{SNP(\text{F9-10 medium dose})} < F_{SNP(\text{F9-10 high dose})} > 0.2$$

Criteria used to identify polymorphisms associated with resistance in our variant analyses

C) Results from the differential expression and transcript polymorphisms analyses.



Venn diagram of SNPs identified in the different strains. Common SNPs between strains are indicated at the intersection of the circles



Correspondence to:

Mathilda Idier
Unité Plantes et Systèmes de culture Horticoles
INRAE Avignon (France)
mathilda.idier@inrae.fr