



Evaluation of the presence of the *bap* gene in *Staphylococcus aureus* isolates recovered from human and animals species.

Eric Vautor, G. Abadie, A. Pont, Régis Thiery

► To cite this version:

Eric Vautor, G. Abadie, A. Pont, Régis Thiery. Evaluation of the presence of the *bap* gene in *Staphylococcus aureus* isolates recovered from human and animals species.. *Veterinary Microbiology*, 2008, 127 (3-4), pp.407-11. 10.1016/j.vetmic.2007.08.018 . hal-00409215

HAL Id: hal-00409215

<https://anses.hal.science/hal-00409215>

Submitted on 16 Nov 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evaluation of the presence of the *bap* gene in *Staphylococcus aureus* isolates recovered from human and animals species

E. VAUTOR*, G ABADIE, A. PONT, R THIERY

AFSSA (French Food Safety Agency), Small Ruminants Pathology Unit, BP111, 06902
Sophia-Antipolis Cedex, France

* Corresponding author. Tel (33) 4 9294 3711; Fax (33) 4 9294 3701.

E-mail address: e.vautor@sophia.afssa.fr

Abstract:

The implication of biofilm in chronic bacterial infection in many species has triggered an increasing interest in the characterization of genes involved in biofilm formation. The *bap* gene is a newly identified gene that encodes the biofilm-associated protein, BAP, which is involved in biofilm formation in *Staphylococcus aureus*. So far the *bap* gene has only been found in a small proportion of *S. aureus* strains from bovine mastitis in Spain. In order to study the presence of the *bap* gene in *S. aureus* isolates obtained from other species and various locations, a collection of 262 isolates was tested by PCR, using published primers and dot-blot. The results indicated that none isolates carried the *bap* gene suggesting that the prevalence of this gene among *S. aureus* isolates should be very low.

Keywords: *Staphylococcus aureus*; *Bap* gene; Biofilm; Epidemiology

1. Introduction

In *Staphylococcus aureus* (*S. aureus*) the implication of biofilm in chronic infections in all animal species have triggered an increasing interest in the characterization of genes

involved in this biofilm formation. For example, the biofilm formation is important for virulence in mastitis (Baselga et al., 1993). A new gene (6,831 nucleotides) involved in biofilm formation (*bap* coding for a biofilm-associated protein, Bap) was identified in a small proportion of *S. aureus* from bovine mastitis (Cucarella et al., 2001). The *bap* protein is a member of proteins playing a role in biofilm formation in many bacteria. They share common structural features as they have a high molecular weight and contain a core domain of tandem repeats. These proteins confer upon bacteria the capacity to form a biofilm and play a relevant role in bacterial infectious process. Some of these proteins are contained occasionally in mobile elements (Lasa and Penades, 2006). In *S. aureus*, the *bap* gene is carried by a putative composite transposon inserted in SaPIbov2, a mobile staphylococcal pathogenicity island. *Bap* orthologue genes have been found in other staphylococcal species including *Staphylococcus epidermidis*, *Staphylococcus chromogenes*, *Staphylococcus xylosus*, *Staphylococcus simulans* and *Staphylococcus hyicus*. However, sequence analyses of the flanking regions revealed that these orthologue *bap* genes of these staphylococcal species were not contained in the SaPIbov2 pathogenicity island (Tormo et al., 2005).

The aim of the present study was to investigate the presence of *bap* gene in various *S. aureus* isolates recovered from human and different animal species. To do this a published PCR method was used (Cucarella et al., 2001) and results were confirmed by dot blot analysis.

2. Materials and methods

2.1. *S. aureus* isolates used in the study

Two hundred and sixty two *S. aureus* isolates associated with different diseases were recovered from various locations in France and different animal species (cows, sheep, goats, pigs, rabbits, poultry, horses, human) (table 1). The cow's isolates were a gift from Dr J.L Martel (AFSSA Lyon). Some sheep isolates were from ML De Buyser (AFSSA Maisons-Alfort). The goat's isolates were partly from Dr P. Mercier (AFSSA Niort). The pig, poultry and

rabbits isolates were from M.H. Băyon-Auboyer (Departemental Laboratory Côtes d'Armor). The horse's isolates were a gift from Dr C. Collobert (AFSSA Dozulé). The human isolates were mainly from Dr H. Carsenti-Dellamonica (Hospital of Archet, Nice, France). All the remaining isolates were from AFSSA Sophia-Antipolis.

2.2. Detection of the *bap* gene by PCR

DNA extraction was performed using the DNeasy[®] Tissue Kit (Qiagen, Courtaboeuf, France) according the manufacturer's instructions with slight modifications. Lysostaphin (Sigma, St Quentin, France) were added (1mg/ml) for enzymatic lysis at 37°C for 2 h.

PCR were performed twice, using a primer pair (*sasp-6m* : 5' CCCTATATCGAAGGTGTAGAATTGCAC 3' and *sasp-7c* : 5' GCTGTTGAAGTTAATACTGTACCTGC 3') as described by Cucarella (Cucarella et al., 2004) to detect the *bap* gene. Amplification was carried out on a Mastercycler[™] (Eppendorf, Hamburg, Germany) with Platinum[®] Taq DNA Polymerase (Invitrogen, Cergy Pontoise, France) under the following conditions: an initial 2 minutes denaturation step at 94°C; followed by 40 cycles each of 30 seconds at 94°C, 30 seconds at 55°C, and 75 seconds at 72°C; and a final step at 72°C for 5 minutes. A 971-bp PCR fragment was expected.

The primer pair (*staur4* : 5' ACGGAGTTACAAAGGACGAC 3' and *staur6* : 5' AGCTCAGCCTTAACGAGTAC 3') was used to target the 23S rDNA as described by Straub (Straub et al., 1999) to confirm the quality of each DNA extract and the absence of PCR inhibitor. The following conditions were used: an initial 5 minutes step at 94°C; followed by 30 cycles each consisting of 30 seconds at 94°C, 30 seconds at 58°C, and 75 seconds at 72°C; and a final step at 72°C for 5 minutes. A 1250-bp PCR fragment was expected.

A *bap* positive control strain V329 (Genbank accession no. AY220730, kindly provided by Dr J.R. Penadés, Spain) was used with each PCR run. Amplification products were electrophoresed in a 1% agarose gel containing ethidium bromide and visualized by transillumination under UV light.

2.3. Dot blotting and hybridisation for the *bap* gene

For the dot blotting, 141 isolates were randomly chosen among the two hundred and sixty two isolates.

The dot blotting technique was described by Planchon et al. (Planchon et al., 2006). Briefly, 50 ng of denatured DNA were spotted onto N + nylon membrane and treated according to the manufacturer's instructions (Amersham Biosciences, Buckinghamshire, England). The PCR product amplified with the primers *sasp-6m* and *sasp-7c* (Cucarella et al., 2004), specific for the *bap* gene (971 bp long) from the V329 *S. aureus* strain, was used as the probe. This PCR product was purified with a QIAquick® PCR purification kit (Qiagen, Courtaboeuf, France), labelled with the DIG-High Prime® system (Roche, Neuilly sur Seine, France). The hybridisations were done in DIG Easy Hyb® solution and the hybridised probe was detected by the Dig colour detection® kit (Roche, Neuilly sur Seine, France) following the manufacturer's instructions. *S. aureus* V329 (Genbank accession no. AY220730) was used as positive control and the strain Mu50 (Genbank accession no. BA000017) as negative control.

3. Results

DNA extracted from 262 *S. aureus* isolates was tested for the presence the *bap* gene by PCR, using the primer pair *sasp-6m* and *sasp-7c*, as indicated in the materials and methods section. Although the positive control strain (V329) showed a band at 971 bp, as expected, none of the tested isolates showed positive results. All isolates were also tested for the presence of the *S. aureus* 23S DNA by using PCR to check for DNA quality, presence of inhibitors of the PCR reactions and specificity. All isolates were found positive, thus eliminating false negative results. An example of some PCR results is shown in figure 1.

Since it cannot be excluded that mutations or deletions could have occurred in the primer pair region of the *bap* gene, 141 isolates were randomly selected and tested by dot

blotting. As illustrated in figure 2, all isolates were also found negative by using this technique, except the *bap* positive control strain V329. Therefore it is likely that the *bap* gene is lacking in all isolates of *S. aureus* tested in this study.

4. Discussion

The gene for the biofilm associated protein (*bap* gene) was not detected in the twenty hundred and sixty two *S. aureus* isolates of this study. These results are in agreement with previous surveys on *S. aureus* of human, bovine, rabbit and pig origins (Arciola et al., 2001; Vasudevan et al., 2003; Vancraeynest et al., 2004; Nitzsche et al., 2007) where the *bap* gene was not found in the *S. aureus* isolates recovered in these animal species of these studies. Our study is the first one with a wide range of *S. aureus* recovered from different animal species to show that the *bap* gene had not spread yet among *S. aureus*.

So far, the *bap* gene has only been found in *S. aureus* obtained from bovine subclinical mastitis in Spain (Cucarella et al., 2001). This gene is also present in other *Staphylococcus* species, including *S. epidermidis*, *S. chromogenes*, *S. xylosus*, *S. simulans* and *S. hyicus* (Tormo et al., 2005; Planchon et al., 2006). But, the *bap* gene is not widely distributed in *S. aureus* isolates despite its presence in the pathogenicity island SaPIbov2, a mobile genetic element. Analysis of the *bap* flanking sequences revealed that *bap* is carried by a transposon-like element. The transposon is inserted in the pathogenic island SaPIbov2 which is mobile without the presence of a helper phage (Penadés, 2006). The ability to produce biofilms, associated with the presence of the *bap* gene has been shown to give growth and persistence advantage to isolates from bovine chronic mastitis (Cucarella et al., 2004). But, *S. aureus* is fully capable of forming biofilm in the absence of *bap* gene as shown for some isolates (Vautor et al., 2006). These isolates were associated with the well-known operon *icaADBC* (manuscript in preparation). As producing biofilm is an advantage for virulence, it was the purpose of this study to look for the *bap* gene in *S. aureus* pathogenic isolates. We propose two hypothesis, to be confirmed, explaining why the *bap* gene had not

spread amongst others *S. aureus* strains: i) the *bap* gene must have been acquired recently by *S. aureus* in SaPIbov2 and consequently the gene has not been horizontally transferred yet ii) horizontal gene transfer is not easy between different *S. aureus* lineages due to their host specificities. The *S. aureus* lineages are supposed to be different between strains recovered from different animal species. The Sau1 type restriction-modification system found in *S. aureus* is one of specific mechanism that controls the ability of mobile genetic element to spread between strains (Waldron and Lindsay, 2006).

In conclusion, this study indicated that none isolates carried the *bap* gene suggesting that the prevalence of this gene among *S. aureus* isolates should be very low. Finally, no evidence of horizontal transfer of the *bap* gene between *S. aureus* recovered from different animal species was found.

Acknowledgements

We thank Dr José Penades (Cardenal Herrera-CEU University and Instituto Valenciano de Investigaciones Agrarias, 46113 Moncada, Valencia, Spain) for helpful discussion.

References

205 Arciola, C.R., Baldassarri, L., Montanaro, L., 2001. Presence of *icaA* and *icaD* genes and
 206 slime production in a collection of staphylococcal strains from catheter-associated
 207 infections. J. Clin. Microbiol. 39, 2151-2156.

208 Baselga, R., Albizu, I., De La Cruz, M., Del Cacho, E., Barberan, M., Amorena, B., 1993.
 209 Phase variation of slime production in *Staphylococcus aureus*: implications in colonization
 210 and virulence. Infect. Immun. 61, 4857-4862.

211 Cucarella, C., Solano, C., Valle, J., Amorena, B., Lasa, I., Penades, J.R., 2001. Bap, a
 212 *Staphylococcus aureus* surface protein involved in biofilm formation. J. Bacteriol. 183,
 213 2888-2896.

214 Cucarella, C., Tormo, M.A., Ubeda, C., Trotonda, M.P., Monzon, M., Peris, C., Amorena, B.,
 215 Lasa, I., Penades, J.R., 2004. Role of biofilm-associated protein bap in the pathogenesis
 216 of bovine *Staphylococcus aureus*. Infect. Immun. 72, 2177-2185.

217 Lasa, I., Penades, J.R., 2006. Bap: a family of surface proteins involved in biofilm formation.
 218 Res. Microbiol. 157, 99-107.

219 Nitzsche, S., Zweifel, C., Stephan, R., 2007. Phenotypic and genotypic traits of
 220 *Staphylococcus aureus* strains isolated from pig carcasses. Vet. Microbiol. 120, 292-299.

221 Penadés, J. R., 2006. Protein-mediated biofilm formation in bacteria: the Bap model.
 222 International Symposium on Staphylococci and Staphylococcal infections, Maastricht, The
 223 Netherlands.[12th, 3-6 September 2006], 63-66.

224 Planchon, S., Gaillard-Martinie, B., Dordet-Frisoni, E., Bellon-Fontaine, M.N., Leroy, S.,
 225 Labadie, J., Hebraud, M., Talon, R., 2006. Formation of biofilm by *Staphylococcus*
 226 *xylosus*. Int. J. Food Microbiol. 109, 88-96.

227 Straub, J.A., Hertel, C., Hammes, W.P., 1999. A 23S rDNA-targeted polymerase chain
 228 reaction-based system for detection of *Staphylococcus aureus* in meat starter cultures
 229 and dairy products. J. Food Prot. 62,1150-1156.

230 Tormo, M.A., Knecht, E., Gotz, F., Lasa, I., Penades, J.R., 2005. Bap-dependent biofilm
 231 formation by pathogenic species of *Staphylococcus*: evidence of horizontal gene transfer?
 232 Microbiology 151, 2465-2475.

233 Vancraeynest, D., Hermans, K., Haesebrouck, F., 2004. Genotypic and phenotypic screening
 234 of high and low virulence *Staphylococcus aureus* isolates from rabbits for biofilm formation
 235 and MSCRAMMs. Vet. Microbiol. 103, 241-247.
 236 Vasudevan, P., Nair, M.K., Annamalai, T., Venkitanarayanan, K.S., 2003. Phenotypic and
 237 genotypic characterization of bovine mastitis isolates of *Staphylococcus aureus* for biofilm
 238 formation. Vet. Microbiol. 92, 179-185.
 239 Vautor, E., Carsenti-Dellamonica, H., Sabah, M., Mancini, G., Pépin, M., Dellamonica, P.,
 240 2006. Characterization of *Staphylococcus aureus* isolates recovered from dairy sheep
 241 farms (*agr* group, adherence, slime, resistance to antibiotics). Small Rumin. Res. In press.
 242 Waldron, D.E., Lindsay, J.A., 2006. Sau1: a novel lineage-specific type I restriction-
 243 modification system that blocks horizontal gene transfer into *Staphylococcus aureus* and
 244 between *S. aureus* isolates of different lineages. J. Bacteriol. 188, 5578-5585.
 245
 246
 247